

Exponent matrices and Frobenius rings

M. A. Dokuchaev, M. V. Kasyanuk,
M. A. Khibina and V. V. Kirichenko

ABSTRACT. We give a survey of results connecting the exponent matrices with Frobenius rings. In particular, we prove that for any permutation $\sigma \in S_n$ there exists a countable set of indecomposable Frobenius semidistributive rings A_n with Nakayama permutation σ .

1. Exponent matrices

Let $M_n(B)$ be the ring of all $n \times n$ matrices over a ring B , a \mathbb{Z} the ring of integers.

An integer matrix $\mathcal{E} = (\alpha_{ij} \in M(\mathbb{Z}))$ is called an exponent matrix if

$$(a) \quad \alpha_{ij} + \alpha_{jk} \geq \alpha_{ik} \text{ and } \alpha_{ii} = 0 \text{ for all } i, j, k = 1, \dots, n;$$

a reduced exponent matrix if

$$(b) \quad \alpha_{ij} + \alpha_{ji} > 0 \text{ for all } i, j = 1, \dots, n; i \neq j.$$

Let

$$\mathcal{E}^{(1)} = (\beta_{ij}), \text{ where } \beta_{ij} = \begin{cases} \alpha_{ij}, & \text{if } i \neq j \\ 1, & \text{if } i = j \end{cases}$$

and

$$\mathcal{E}^{(2)} = (\gamma_{ij}), \text{ where } \gamma_{ij} = \min_{1 \leq k \leq n} (\beta_{ik} + \beta_{kj}).$$

2010 MSC: 16P40, 16P20.

Key words and phrases: exponent matrix, Frobenius ring, distributive module, quiver of semiperfect ring.

Consider the matrix $[Q] = \mathcal{E}^2 - \mathcal{E}^1 = (q_{ij})$. Let $\beta_{ik} + \beta_{kj}$ ($i \neq j$) be an integer. If $k = i$, then $\beta_{ii} + \beta_{ij} = \beta_{ii} + \alpha_{ij} = \alpha_{ij} + 1$.

Therefore $\min_{1 \leq k \leq n} (\beta_{ik} + \beta_{kj}) \leq \alpha_{ij} + 1$ and $[Q]$ is $(0, 1)$ -matrix and it is the adjacency matrix of a simply laced quiver.

Definition 1.1. A reduced exponent matrix $\mathcal{E} = (\alpha_{ij}) \in M_n(\mathbb{Z})$ is called Gorenstein if there exists a permutation σ of $\{1, 2, \dots, n\}$ such that $\alpha_{ik} + \alpha_{k\sigma(i)} = \alpha_{i\sigma(i)}$ for $i, k = 1, \dots, n$.

The permutation σ is denoted by $\sigma(\mathcal{E})$. Obviously, $\sigma(\mathcal{E})$ has no cycles of length 1.

Gorenstein matrices are closely related to the semiperfect semidistributive rings A with non-zero Jacobson radical and $inj.dim_A A_A = 1$ [1].

Consider a reduced symmetric exponent matrix $\mathcal{E} = (\alpha_{ij}) \in M_n(\mathbb{Z})$. It means $\alpha_{ij} + \alpha_{ji} > 0$ for $i \neq j$, i.e., $2\alpha_{ij} > 0$ for $i \neq j$. Thus, all nondiagonal elements of the reduced symmetric matrix are positive, and it defines a finite metric space.

Definition 1.2. A metric space is a pair (M, d) , where M is a set and $d : M \times M \rightarrow [0, \infty)$ is a metric, satisfying the following axioms:

- (1) $d(x, y) = 0$, if and only if $x = y$;
- (2) $d(x, y) = d(y, x)$ for all $x, y \in M$;
- (3) $d(x, y) + d(y, x) \geq d(x, z)$ for all $x, y, z \in M$.

If M is finite, then M is a finite metric space. Denote $m_{ij} = d(i, j)$. The matrix $D(M) = (m_{ij}) \in M_n(\mathbb{R})$ is called the distance matrix of M .

Now we give the examples of Gorenstein matrices.

Example 1.1. The following exponent matrix

$$T_{n,\alpha} = \begin{pmatrix} 0 & 0 & \dots & \dots & \dots & 0 \\ \alpha & 0 & 0 & \dots & \dots & 0 \\ \alpha & \ddots & \ddots & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \alpha & & \ddots & \alpha & 0 & 0 \\ \alpha & \dots & \dots & \alpha & \alpha & 0 \end{pmatrix}$$

is Gorenstein with $\sigma(T_{n,\alpha}) = (12 \dots n)$.

Example 1.2. The Cayley table of the Klein four-group $(2) \times (2)$ can be written in the following form:

$$K(4) = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 1 & 0 & 3 & 2 \\ 2 & 3 & 0 & 1 \\ 3 & 2 & 1 & 0 \end{pmatrix}.$$

$K(4)$ is Gorenstein with $\sigma(K(4)) = (14)(23)$.

Example 1.3. The Cayley table of the elementary Abelian n_2 -group $(2) \times (2) \times (2)$ is as following

$$K(8) = \begin{pmatrix} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 0 & 3 & 2 & 5 & 4 & 7 & 6 \\ 2 & 3 & 0 & 1 & 6 & 7 & 4 & 5 \\ 3 & 2 & 1 & 0 & 7 & 6 & 5 & 4 \\ 4 & 5 & 6 & 7 & 0 & 1 & 2 & 3 \\ 5 & 4 & 7 & 6 & 1 & 0 & 3 & 2 \\ 6 & 7 & 4 & 5 & 2 & 3 & 0 & 1 \\ 7 & 6 & 5 & 4 & 3 & 2 & 1 & 0 \end{pmatrix}.$$

$K(8)$ is Gorenstein with $\sigma(K(8)) = (18)(27)(36)(45)$.

The notion of a Latin square was introduced by L. Euler at the end of the XVIII century (see [2] for details).

Definition 1.3. A Latin square \mathcal{L}_n of order n is a square $n \times n$ matrix of order n , such that its rows and columns are permutations of some set $S = \{s_1, \dots, s_n\}$.

In what follows we shall take $S = \{0, 1, \dots, n-1\}$. So, $\mathcal{L}_n = (\alpha_{ij})$, where $\alpha_{ij} \in \{0, 1, \dots, n-1\}$.

We say that a Latin square of order n is normalized if its first row is $(0, 1, \dots, n-1)$ and the first column is $(0, 1, \dots, n-1)^T$, where T (as exponent index) means transposition. The normalized Latin squares are also called “reduced Latin squares” or “Latin squares of standard form”.

We use the following notations:

$$\Gamma_0 = (0), \quad \Gamma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \Gamma_2 = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 1 & 0 & 3 & 2 \\ 2 & 3 & 0 & 1 \\ 3 & 2 & 1 & 0 \end{pmatrix},$$

$$U_n = \begin{pmatrix} 1 & 1 & \dots & 1 \\ 1 & 1 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & 1 \end{pmatrix} \in M_n(\mathbb{Z}), \quad X_{k-1} = 2^{k-1}U_{2^{k-1}};$$

$$\Gamma_k = \begin{pmatrix} \Gamma_{k-1} & \Gamma_{k-1} + X_{k-1} \\ \Gamma_{k-1} + X_{k-1} & \Gamma_{k-1} \end{pmatrix} \text{ for } k = 1, 2, \dots$$

The matrix $\Gamma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ is the Cayley table of the cyclic group G_1 of order 2 and is a Gorenstein matrix with permutation $\sigma(\Gamma_1) = (12)$.

Proposition 1.4 ([3, §7.6]). Γ_k is an exponent matrix for any natural number k .

Proposition 1.5 ([3, §7.6]). Γ_k is the Cayley table of the elementary Abelian group G_k of order 2^k .

Proposition 1.6 ([3, §7.6]). The matrix Γ_k is Gorenstein with permutation

$$\sigma(\Gamma_k) = \begin{pmatrix} 1 & 2 & 3 & \dots & 2^k - 1 & 2^k \\ 2^k & 2^k - 1 & 2^k - 2 & \dots & 2 & 1 \end{pmatrix}.$$

Theorem 1.7 ([4] and [3, §7.6]). Suppose that a Latin square \mathcal{L}_n with first row and first column $(0\ 1\ \dots\ n - 1)$ is an exponent matrix. Then $n = 2^k$ and $\mathcal{L}_n = \Gamma_k$ is the Cayley table of the direct product of k copies of the cyclic group of order 2.

Conversely, the Cayley table Γ_k of the elementary Abelian group

$$G_k = \mathbb{Z}/(2) \times \dots \times \mathbb{Z}/(2) = (2) \times \dots \times (2)$$

(k factors) of order 2^k is a Latin square and a Gorenstein symmetric matrix with the first row $(0, 1, \dots, 2^k - 1)$ and permutation

$$\sigma(\Gamma_k) = \begin{pmatrix} 1 & 2 & 3 & \dots & 2^k - 1 & 2^k \\ 2^k & 2^k - 1 & 2^k - 2 & \dots & 2 & 1 \end{pmatrix}.$$

Now we consider the case when a Latin square \mathcal{L}_n with first row and first column $(0\ 1\ \dots\ n - 1)$ is a distance matrix $D = D(M)$ of a finite metric space $M = \{m_1, \dots, m_n\}$. Obviously, if $\mathcal{L}_n = D(M)$ then \mathcal{L}_n is an exponent matrix. So we obtain the following theorem.

Theorem 1.8. *Suppose that a Latin square \mathcal{L}_n with first row and first column $(0, 1, \dots, n-1)$ is a distance matrix $D = D(M)$ of a finite metric space $M = \{m_1, \dots, m_n\}$. Then $n = 2^k$ and $\mathcal{L}_n = \Gamma_k$ is the Cayley table of the direct product of k copies of the cyclic group of order 2.*

Conversely, the Cayley table Γ_k of the elementary Abelian group

$$G_k = \mathbb{Z}/(2) \times \dots \times \mathbb{Z}/(2) = (2) \times \dots \times (2)$$

(k factors) of order 2^k is a Latin square and the distance matrix $D = D(M)$ of a finite metric space with 2^k elements.

2. Semiperfect rings

Recall that a ring A is called local, if it has a unique maximal right ideal \mathfrak{M} .

An idempotent $e \in A$ is called local if the ring eAe is local. Clearly, a local idempotent is always a primitive idempotent. We shall say that idempotents can be lifted modulo an ideal \mathcal{I} of a ring A if from the fact that $g^2 - g \in \mathcal{I}$, where $g \in A$, it follows that there exists an idempotent $e^2 = e \in A$ such that $e - g \in \mathcal{I}$ (or that the same $g - e \in \mathcal{I}$).

Show that in a local ring an idempotents can be lifted modulo a unique maximal ideal R . Consider $g^2 - g \in R$. So $g(g - 1) \in R$. If $g \in R$, then $g - 0 = g \in R$. If $g \notin R$, then $g - 1 \in R$.

Proposition 2.1. *Idempotents can be lifted modulo any nil-ideal I of a ring A .*

Proof. Let $g^2 - g \in I$. Set $r = g^2 - g$, $g_1 = g + r - 2gr$. Obviously, $rg = g^3 - g^2 = g(g^2 - g) = gr$. Calculating $g_1^2 - g_1$, we obtain $g_1^2 = (g + r - 2gr)^2 = (g+r)^2 + 4g^2r^2 - 4gr(g+r) = g^2 + 2gr + r^2 + 4g^2r^2 - 4g^2r - 4gr^2$. Therefore, $g_1^2 - g_1 = 2gr + r^2 + 4g^2r^2 - 4g^2r - 4gr^2 + 2gr = r^2 + 4g^2r^2 + 4gr - 4g^2r - 4gr^3 = r^2 - 4r^3 - 4r^2 = r^2(4r - 3)$.

Let $r_1 = r^2(4r - 3) \in I$ and $g_1 = g_1 + r_1 - 2g_1r_1$. We obtain that $r_2 = g_1^2 - g_1 = r_1^2(4r_1 - 3)$, i.e., in the expression of r_2 the element r^4 enters a factor. Since $r^k = 0$ for some integer $k > 0$, we obtain that $r_n = 0$ for some n , i.e., $g_n^2 = g_n$. We have that $g_1 - g \in I$ and $g_i - g_{i-1} \in I$ for $i = 1, 2, \dots, n$ and we obtain $g_n = e$ is idempotent and $g - e \in I$ \square

Theorem 2.2 (Hopkins-Levitsky Theorem). *The Jacobson radical R of a right Artinian ring A is nilpotent [5], [6].*

Therefore, any right Artinian ring is semiperfect.

Lemma 2.3. *Let a ring A have two decompositions into a direct sum of right ideals: $A = e_1A \oplus \dots \oplus e_nA = f_1A \oplus \dots \oplus f_nA$ (where $1 = e_1 + \dots + e_n = f_1 + \dots + f_n$ are two decompositions of $1 \in A$ into a sum of pairwise orthogonal idempotents), and suppose $e_iA \simeq f_iA$ ($i = 1, \dots, n$), after renumbering if necessary. Then there is an invertible element $a \in A$ such that $f_i = ae_i a^{-1}$.*

Proof. Let $\varphi : e_iA \rightarrow f_iA$ be the isomorphism. Then $\varphi(e_i a) = \varphi(e_i) a$ and $\varphi(e_i) = \varphi(e_i^2) = \varphi(e_i) e_i$. We have $\varphi(e_i) = f_i a$. So $\varphi(e_i) = f_i a e_i$ and endomorphism φ is realized by multiplication on the left side by element $a_i = f_i a e_i$. Then $f_i a_i = a_i e_i = a_i$. Consider the elements $b_i \in e_i a f_i$ realizing the inverse isomorphisms. We set $b = b_1 + \dots + b_n$ and $a = a_1 + \dots + a_n$. Obviously, $e_i b = b_i = b f_i$. Moreover, $a_i b_i = f_i$, $b_i a_i = e_i$. Clearly, $ab = \sum_{i=1}^n a_i b_i = \sum_{i=1}^n f_i = 1$ and $ba = \sum_{i=1}^n b_i a_i = \sum_{i=1}^n e_i = 1$, i.e., $b = a^{-1}$. Since $ae_i = f_i a$, we have $f_i = ae_i a^{-1}$. □

We will use the following notations and definitions.

Let M be a module over a ring A with the Jacobson radical R . Denote by $\text{rad } M$ the intersection of all its maximal submodules. By convention, if M does not have maximal submodules we define $\text{rad } M = M$. This submodule is called the radical of the module M .

Recall that a right ideal \mathfrak{M} in a ring A is called maximal in A if there is no right ideal J , different from \mathfrak{M} and A such that $\mathfrak{M} \subset J \subset A$. In nonzero ring with identity always there exist maximal proper right ideals.

The following proposition is often useful.

Proposition 2.4. *Let M be a right A -module, and let R be the Jacobson radical of the ring A . Then $MR \subset \text{rad } M$.*

Theorem 2.5. *A ring A is semiperfect if and only if it can be decomposed into a direct sum of right ideals each of which has exactly one maximal submodule.*

Proof. Let $\bar{A} = A/R = \bar{e}_1 \bar{A} \oplus \dots \oplus \bar{e}_n \bar{A}$ be a decomposition of \bar{A} into a direct sum of simple \bar{A} -modules. Write $\bar{e}_i \bar{A} = U_i$. Since A is semiperfect, for each idempotent \bar{e}_i there is an idempotent e_i such that $e_i + R = \bar{e}_i$. Let $P_i = e_i A$. Obviously, $P_i R = e_i A R \subset R$. Therefore $P_i \cap R = P_i R$. Consequently, $P_i + R/R \simeq P_i/P_i R \simeq U_i$ and every module P_i has exactly one maximal submodule $P_i R$. Let $P = \bigoplus_{i=1}^n P_i$. Obviously, there is an epimorphism $\varphi : P \rightarrow \bar{A}$. Denote by π the natural projection $A \rightarrow \bar{A}$. Since

the module P is projective, there exists a homomorphism $\psi : P \rightarrow A$ such that $\pi\psi = \varphi$. Obviously, $Im\psi + R = A$. By Nakayama's lemma $Im\psi = A$. Since the module A is projective, we have $P \simeq Im\psi \oplus Ker\psi = A \oplus Ker\psi$. We want show that $X = Ker\psi = 0$. Consider P/PR . Then $P/PR \simeq \bar{A}$ and, on the other hand $P/PR = \bar{A} \oplus X/XR$. By the Krull-Schmidt theorem for semisimple modules we obtain that the module X/XR is equals to zero. By Nakayama's lemma $X = 0$, i.e., X is finitely generated as the image P . Therefore $Ker\psi = 0$ and A decomposes into a direct sum of right ideals $\psi(e_i A)$, each of which has exactly one maximal submodule.

Conversely, let $A_A = P_1 \oplus \dots \oplus P_n$ be a decomposition of right regular A -module A into a direct sum of right A -modules, each of which has exactly one maximal submodule $rad P_i$ ($i = 1, \dots, n$). Consequently, $R = rad P_1 \oplus \dots \oplus rad P_n$ and $\bar{A} = A/R$ is a right semisimple ring. Let $1 = f_1 + \dots + f_n$ be a decomposition of the identity of the ring A into a sum of pairwise orthogonal idempotents such that $P_i = f_i A$ ($i = 1, \dots, n$). \square

We shall show that for any idempotent $\bar{e}^2 = \bar{e} \in \bar{A}$ there is an idempotent $e \in A$ such that $e + R = \bar{e}$. The right ideal $\bar{e}\bar{A}$ is semisimple as \bar{A} -submodule of the semisimple module \bar{A} . Therefore there is a decomposition of $\bar{1} \in \bar{A}$ into a sum of pairwise orthogonal idempotents $\bar{1} = \bar{e}_1 + \dots + \bar{e}_s + \dots + \bar{e}_n$ such that $\bar{e} = \bar{e}_1 + \dots + \bar{e}_s$ and all modules $\bar{e}_i \bar{A}$ are simple. On the other hand, let $\bar{1} = \bar{f}_1 + \dots + \bar{f}_n$ be a decomposition of $\bar{1} \in \bar{A}$ into a sum of pairwise orthogonal idempotents such that the modules $\bar{f}_i \bar{A}$ are simple. By Krull-Schmidt theorem for semisimple modules for an appropriate renumeration we have $\bar{e}_i \bar{A} \simeq \bar{f}_i \bar{A}$ ($i = 1, \dots, n$). By lemma 2.3 there exists an element $\bar{a} \in \bar{A}$ such that $\bar{e}_i = \bar{a}^{-1} \bar{f}_i \bar{a}$ ($i = 1, \dots, n$). Let \bar{a} be the image of a and \bar{a}^{-1} be the image of $b = a^{-1}$. Obviously, $ab = 1 + r$, where $r \in R$. Since $(1+r)x = 1$ we have $x = 1 - r_1$, where $r_1 \in R$. Therefore, $b(1 - r_1) = bx = 1$. So, $bx = a^{-1}$. $bx = b - br_1$ and the image of the element a^{-1} under the epimorphism π coincides with \bar{a}^{-1} . Then $\pi(e) = \sum \pi(a^{-1} f_i a) = \sum \bar{a}^{-1} \bar{f}_i \bar{a} = \bar{e}$. The theorem is proved.

Theorem 2.6 (B. J. Mueller). *A ring A is semiperfect if and only if $1 \in A$ can be decomposed into a sum of a finite number of pairwise orthogonal local idempotents.*

Proof. Let a ring A be semiperfect. By previous theorem $A = e_1 A \oplus \dots \oplus e_n A$, where e_1, \dots, e_n are idempotents and each right ideal $P_i = e_i A$ ($i = 1, \dots, n$) has exactly one maximal submodule. Then $Hom(P_i, P_i) \simeq e_i A e_i$ and for any $\psi : P_i \rightarrow P_i$ either $Im\psi = P_i$ or $Im\psi \subset P_i R$. In the first

case, since $Im \psi \simeq P_i$ is projective we have $P_i \simeq Im \psi \oplus Ker \psi$, which implies $Ker \psi = 0$ and so ψ is an automorphism. In the second case, ψ is noninvertible element and, obviously, all noninvertible elements form an ideal. Therefore, the ring $e_i A e_i$ is local for $i = 1, \dots, n$.

Conversely, let $\pi : A \rightarrow \bar{A}$ be the natural projection of the ring A on the ring $\bar{A} = A/R$. (R is the Jacobson radical of the ring A). Let $\pi(a) = \bar{a}$ and e be a local idempotent of the ring A . We shall show that the module $\pi(eA) = \bar{e}\bar{A}$ is simple. We may consider that the ring A is not local (a local ring is semiperfect). Consider $(\bar{1} - \bar{e})\bar{A}$. Since it is a proper right ideal in the ring \bar{A} , it is contained in a maximal right ideal \tilde{I} of the ring \bar{A} . We shall show that $\bar{e}\bar{A} \cap \tilde{I} = 0$. If $(\bar{e}\bar{A} \cap \tilde{I}) \neq 0$, then $(\bar{e}\bar{A} \cap \tilde{I})^2 \neq 0$. The ring A/R is semiprimitive ring and therefore it has no nilpotent right ideals. Then there is an element $\bar{e}\bar{a} \in \tilde{I}$ and $\bar{e}\bar{a}\bar{e}\bar{a} \neq 0$. So $\bar{e}\bar{a}\bar{e} \neq 0$. Since eAe is a local ring and $rad(eAe) = eRe$, we conclude that the ring $\bar{e}\bar{A}\bar{e}$ is a division ring. Therefore, there is an element $\bar{e}\bar{x}\bar{e} \in \bar{e}\bar{A}\bar{e}$ such that $\bar{e}\bar{a}\bar{e}\bar{x}\bar{e} = \bar{e}$ and $\bar{e} \in \tilde{I}$. Thus $\bar{1} \in \tilde{I}$, a contradiction. We obtain, that $\bar{e}\bar{A} \cap \tilde{I} = 0$ and $\bar{A} = \bar{e}\bar{A} \oplus \tilde{I}$. Since \tilde{I} is the maximal ideal in the ring A , the module $\bar{e}\bar{A}$ is simple. The theorem is proved. \square

3. Distributive modules

Recall that a module M is called distributive if for all submodules K, L, N of M we have $K \cap (L + N) = K \cap L + K \cap N$. Clearly, a submodule and a quotient module of a distributive module is distributive. A module is called semidistributive if it is a direct sum of distributive modules. A ring A is called right (left) semidistributive if the right (left) regular module A_A (${}_A A$) is semidistributive. A right and left semidistributive ring is called semidistributive. Obviously, every uniserial module is a distributive module and every serial module is a semidistributive module.

Example 3.1. Let D be a division ring and

$$A = \left\{ \left(\begin{array}{ccc} d_{11} & d_{12} & d_{13} \\ 0 & d_{22} & 0 \\ 0 & 0 & d_{33} \end{array} \right) \mid d_{ij} \in D \right\}.$$

Obviously, A is the semidistributive ring, which is left serial, but not right serial.

We write *SPSD*-ring for semiperfect semidistributive ring.

4. Semiprime right Noetherian *SPSD*-rings

Definition 4.1. A ring A is called *semimaximal* if it is a semiperfect semiprime right Noetherian ring such that for each local idempotent $e \in A$ the ring eAe is a discrete valuation ring (not necessary commutative), i.e., all principal endomorphism rings of A are discrete valuation rings.

The following is a decomposition theorem for semiprime right Noetherian *SPSD*-rings.

Theorem 4.2. The following conditions for a semiperfect semiprime right Noetherian ring A are equivalent:

- (a) the ring A is semidistributive;
- (b) the ring A is a direct product of a semisimple Artinian ring and a semimaximal rings.

Theorem 4.3. Each semimaximal ring is isomorphic to a finite direct product of prime rings of the following form:

$$A = \begin{pmatrix} \mathcal{O} & \pi^{\alpha_{12}}\mathcal{O} & \dots & \pi^{\alpha_{1n}}\mathcal{O} \\ \pi^{\alpha_{21}}\mathcal{O} & \mathcal{O} & \dots & \pi^{\alpha_{2n}}\mathcal{O} \\ \dots & \dots & \dots & \dots \\ \pi^{\alpha_{n1}}\mathcal{O} & \pi^{\alpha_{n2}}\mathcal{O} & \dots & \mathcal{O} \end{pmatrix},$$

where $n \geq 1$, \mathcal{O} is a discrete valuation ring with a prime element π , and the α_{ij} are integers such that $\alpha_{ij} + \alpha_{jk} \geq \alpha_{ik}$ for all i, j, k ($\alpha_{ii} = 0$ for any i).

Definition 4.4. A ring A is called a *tiled order* if it is a prime Noetherian *SPSD*-ring with nonzero Jacobson radical.

Our definition of a tiled order is a generalization of the definition of a tiled order over a discrete valuation ring in the sense V. A. Jategaonkar [7], [8].

We'll use the following notation $A = \{\mathcal{O}, \mathcal{E}(A)\}$, where $\mathcal{E}(A) = (\alpha_{ij})$ is the exponent matrix of a ring A , i.e., $A = \sum_{i,j=1}^n e_{ij}\pi^{\alpha_{ij}}\mathcal{O}$, where the e_{ij} are the matrix units. If a tiled order A is reduced, i.e., A/R is a direct product of division rings, then the matrix $\mathcal{E}(A)$ also reduced, i.e., $\alpha_{ij} + \alpha_{ji} > 0$ for $i, j = 1, \dots, n$ ($i \neq j$).

It is well-known the following proposition.

Proposition 4.5 ([9, vol II, proposition 22.1.A]). *Let P be nonzero projective A -module and $rad A$ be the Jacobson radical of A . Then $rad P = Prad A \neq P$.*

Using this proposition and Muller’s theorem, we obtain the description of projective modules over a semiperfect ring.

Theorem 4.6. *Any indecomposable projective module over a semiperfect ring A is finitely generated, and has exactly one maximal submodule. There is one-to-one correspondence between pairwise nonisomorphic indecomposable projective A -modules P_1, \dots, P_s and mutually nonisomorphic simple A -modules? which are given by the following correspondence: $P_i \rightarrow P_i/P_iR = U_i$ and $U_i \rightarrow P(U_i)$.*

Remark. A submodule N of a module M is called small if the equality $N + X = M$ implies $X = M$ for any submodule X of M .

A projective module P is called a projective cover of M and it is denoted by $P(M)$ if there is an epimorphism $\varphi : P \rightarrow M$ such that $Ker \varphi$ is a small submodule in P . The following is a method of constructing a projective cover for an arbitrary finitely generated module M over a semiperfect ring A . Let M - be a finitely generated A -module. Clearly, M/MR is a module over the semisimple Artinian ring $\bar{A} = A/R$. Therefore $M/MR \simeq \bigoplus_{j=1}^s U_j^{m_j}$, where U_1, \dots, U_s are mutually nonisomorphic simple \bar{A} -modules. Lifting the idempotents we obtain that $U_j = e_jA/e_jR$, where $e_j^2 = e_j$. Denote $P_j = e_jA$. Then the projective cover $P(M)$ of M is $P(M) = \bigoplus_{j=1}^s P_j^{m_j}$.

5. Quiver of a semiperfect ring

Let $A_A = P_1^{n_1} \oplus \dots \oplus P_s^{n_s}$ be the decomposition of a semiperfect ring A into a direct sum of pairwise nonisomorphic indecomposable projectives. Let $P = P_1 \oplus \dots \oplus P_s$ and $B = End_A P$. By the Morita theorem, the category of right A -modules is equivalent to the category of right B -modules. Obviously, $B/rad B$ is a direct product of division rings. The ring B is called the basic ring of the ring A .

Let A be a semiperfect right Noetherian ring and $A_A = P_1^{n_1} \oplus \dots \oplus P_s^{n_s}$ be the decomposition A as above. Let P_iR is the unique maximal

submodule of P_i . Consider the projective cover $P(P_iR)$ of P_iR . Let $P(P_iR) = \bigoplus_{j=1}^s P_j^{t_{ij}}$ ($i = 1, \dots, s$). We assign to the modules P_1, \dots, P_s points $1, \dots, s$ on the plane and join point i with j by t_{ij} arrows. The so constructed graph is called the right quiver of the semiperfect right Noetherian ring A and will be denoted by $Q(A)$.

Analogously, one can define the left quiver $Q'(A)$ of a left Noetherian semiperfect ring.

Note, that the quiver of a semiperfect right Noetherian ring does not change by switching to its basic ring and $Q(A) = Q(A/R^2)$.

Definition 5.1. *Let A be a semiperfect ring such that A/R^2 is a right Artinian ring. The quiver of the ring A/R^2 is the quiver of the ring A and is denoted by $Q(A)$.*

The following proposition gives the description of the Jacobson radical of semiperfect ring A (see [10, Chapter 11]).

Proposition 5.2. *Let $A = P_1^{n_1} \oplus \dots \oplus P_s^{n_s}$ be the decomposition of a semiperfect ring A into a direct sum of principal right A -modules and let $1 = f_1 + \dots + f_s$ be a corresponding decomposition of the identity of A into a sum of pairwise orthogonal idempotents, i.e., $f_i A = P_i^{n_i}$. Then the Jacobson radical of the ring A has a two-sided Peirce decomposition of the following form:*

$$R = \begin{pmatrix} R_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & R_{22} & \dots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{n1} & A_{n2} & \dots & R_{nn} \end{pmatrix},$$

where $R_{ii} = \text{rad}(f_i A f_i)$, $A_{ij} = f_i A f_j$ for $i, j = 1, \dots, n$.

Lemma 5.3 (Annihilation lemma). *Let $1 = f_1 + \dots + f_s$ be a canonical decomposition of $1 \in A$. For every simple right A -module U_i and for each f_j we have $U_i f_j = \delta_{ij} U_i$, $i, j = 1, \dots, s$. Similarly, for every simple left A -module V_i and for each f_j , $f_j V_i = \delta_{ij} V_i$, $i, j = 1, \dots, s$.*

Lemma 5.4 (Q -Lemma). *The simple module U_k (resp. V_k) appears in the direct sum decomposition of the module $e_i R / e_i R^2$ (resp. $R e_i / R^2 e_i$) if and only if $e_i R^2 e_k$ (resp. $e_k R^2 e_i$) is strictly contained in $e_i R e_k$ (resp. $e_k R e_i$).*

6. Quasi-Frobenius rings

Definition 6.1. *Let M be a right A -module. The socle of M ($\text{soc } M$) is the sum of all simple right submodules of M . If there are no such submodules, then $\text{Soc } M = 0$.*

If $M = A_A$, then $\text{soc } A_A$ is the sum of all minimal right ideals of A and it is a right ideal of A . If \mathcal{J} a minimal right ideal in A , then for any $x \in A$ either $x\mathcal{J} = 0$ or $x\mathcal{J}$ is a minimal right ideal in A . Therefore, $\text{soc } A_A$ is a two-sided ideal in A . Analogously we can consider $\text{soc}_A A$. However $\text{soc } A_A \neq \text{soc}_A A$ in general. Let $A = T_2(k) = \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & \gamma \end{pmatrix} \mid \alpha, \beta, \gamma \in k \right\}$, where k is a field. Obviously, $\text{soc } A_A = \left\{ \begin{pmatrix} 0 & \beta \\ 0 & \gamma \end{pmatrix} \right\}$ and $\text{soc}_A A = \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & 0 \end{pmatrix} \right\}$. For semisimple module M we have $\text{soc } M = M$.

Let A be an Artinian ring and $A_A = P_1^{n_1} \oplus \dots \oplus P_s^{n_s}$ (resp. ${}_A A = Q_1^{n_1} \oplus \dots \oplus Q_s^{n_s}$) be the canonical decomposition of A into a direct product of right (left) principal modules.

Let M be a right A -module and N be a left A -module. We set $\text{top } M = M/MR$ and $\text{top } N = N/RN$. Now we give a definition of a Nakayama permutation.

Definition 6.2. *We say that a two-sided Artinian ring A admits a Nakayama permutation $\nu(A) : i \rightarrow \nu(i)$ of $\{1, \dots, s\}$ if the following conditions are satisfied*

- (1) $\text{soc } P_k = \text{top } P_{\nu(k)}$
- (2) $\text{soc } Q_{\nu(k)} = \text{top } Q_k$

Definition 6.3. *A two-sided Artinian ring A is called quasi-Frobenius, if A admits a Nakayama permutation $\nu(A)$ of $\{1, \dots, s\}$. A quasi-Frobenius ring is called Frobenius if $n_{\nu(i)} = n_i$ for all $i = 1, \dots, s$.*

The following theorem is a variant of the Morita theorem for semiperfect rings.

Theorem 6.4. *Let A be a semiperfect ring and $A_A = P_1^{n_1} \oplus \dots \oplus P_s^{n_s}$ be a decomposition of A_A into a direct sum of pairwise nonisomorphic right ideals. Let $B = \text{End}(P_1 \oplus \dots \oplus P_s)$ be the ring of endomorphisms of the module $P = P_1 + \dots + P_s$. Then the categories of A -modules and B -modules are equivalent.*

Denote by $\text{rad } B$ the Jacobson radical of the ring B . Obviously, $B/\text{rad } B$ is a direct product of skew-fields. The ring B is called a basic ring of the ring A . If a ring A is quasi-Frobenius, B is Frobenius.

Definition 6.5. *An indecomposable projective right module over a semi-perfect ring A is called principal right module. Analogously, a principal left module can be defined.*

7. Examples

7.1.

Definition 7.1. *An A -lattice M is said to be relatively injective if $M \simeq_A P^*$, where ${}_A P$ is a finitely generated projective left A -module. An A -lattice M is called completely decomposable if it is a direct sum of irreducible A -lattices.*

Denote by Δ the completely decomposable left A -lattice A_A^* .

Let $A = \{\mathcal{O}, \mathcal{E}(A)\}$, where $\mathcal{E}(A) = (\alpha_{ij})$ be a reduced exponent matrix.

Let

$$\mathcal{E}(A) = \begin{pmatrix} 0 & \alpha_{12} & \dots & \alpha_{1n} \\ \alpha_{21} & 0 & \dots & \alpha_{2n} \\ \dots & \dots & \dots & \dots \\ \alpha_{n1} & \alpha_{n2} & \dots & 0 \end{pmatrix}.$$

Lemma 7.2. *A completely decomposable left A -lattice Δ is complete decomposable right A -lattice, and*

$$\mathcal{E}(\Delta) = \begin{pmatrix} 0 & -\alpha_{21} & \dots & -\alpha_{n1} \\ -\alpha_{12} & 0 & \dots & -\alpha_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ -\alpha_{1n} & -\alpha_{2n} & \dots & 0 \end{pmatrix}.$$

Let us show that the k -th row $(-\alpha_{1k}, -\alpha_{2k}, \dots, -\alpha_{nk})$ of the matrix $\mathcal{E}(\Delta)$ defines an irreducible right A -lattice. Let $\beta_i = -\alpha_{ik}$. We can rewrite the inequality $\alpha_{ij} + \alpha_{jk} \geq \alpha_{ik}$ in the form $-\alpha_{ik} + \alpha_{ij} \geq -\alpha_{jk}$, i.e., $\beta_i + \alpha_{ij} \geq \beta_j$, which implies the assertion of the lemma.

Corollary 7.3. *A fractional ideal Δ is a relatively injective right and a relatively injective left A -lattice.*

Let A be a reduced tiled order and $R = \text{rad } A$. Then

$$\mathcal{E}(R) = \begin{pmatrix} 1 & \alpha_{12} & \dots & \alpha_{1n} \\ \alpha_{21} & 1 & \dots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{n1} & \alpha_{n2} & \dots & 1 \end{pmatrix}$$

and

$$\mathcal{E}(R_A^*) = \mathcal{E}({}_A R^*) = \begin{pmatrix} -1 & -\alpha_{12} & \dots & -\alpha_{1n} \\ -\alpha_{21} & -1 & \dots & -\alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ -\alpha_{n1} & -\alpha_{n2} & \dots & 1 \end{pmatrix}$$

write $X = {}_A R^*$, $\Delta = (A_A)^*$.

It is true the following lemma.

Lemma 7.4. *For $i = 1, \dots, n$ we have that $e_{ii}X(Xe_{ii})$ is the unique minimal overmodule of $e_{ii}A(\Delta e_{ii})$ and $e_{ii}X/e_{ii}\Delta = U_i$, $Xe_{ii}/\Delta e_{ii} = V_i$, where U_i is simple right A -module and V_i is a simple left A -module.*

Lemma 7.5. *Every relatively injective irreducible A -lattice Q has only one minimal overmodule. Let Q_1 and Q_2 be relatively injective irreducible A -lattices and let $X_1 \supset Q$ and $X_2 \supset Q_2$ be the unique minimal overmodules of Q_1 and Q_2 respectively. Then the simple A -modules X_1/Q_1 and X_2/Q_2 are isomorphic if and only if $Q_1 \simeq Q_2$.*

Proposition 7.6. *An irreducible A -lattice is relatively injective if and only if it has exactly one minimal overmodule.*

Consider the tiled order

$$H_n(\mathcal{O}) = \begin{pmatrix} \mathcal{O} & \mathcal{O} & \dots & \mathcal{O} \\ \pi\mathcal{O} & \mathcal{O} & \dots & \mathcal{O} \\ \dots & \dots & \dots & \dots \\ \pi\mathcal{O} & \pi\mathcal{O} & \dots & \mathcal{O} \end{pmatrix},$$

i.e., $H = H_n(\mathcal{O}) = \{\mathcal{O}, \mathcal{E}(H_n(\mathcal{O}))\}$, where

$$\mathcal{E}(H_n(\mathcal{O})) = \begin{pmatrix} 0 & 0 & \dots & 0 \\ 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 1 & 1 & \dots & 0 \end{pmatrix},$$

i.e., $\alpha_{ij} = 0$ for $i \leq j$ and $\alpha_{ij} = 1$ for $i > j$; $i, j = 1, \dots, n$.

Let $\Delta = H_H^*$. Obviously,

$$\mathcal{E}(\Delta) = \begin{pmatrix} 0 & -1 & \dots & -1 \\ 0 & 0 & \dots & -1 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}.$$

Let $M_n(\mathbb{Z})$ be the ring of $n \times n$ -matrices over the ring of integers

\mathbb{Z} . Let $U_n = \begin{pmatrix} 1 & \dots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \dots & 1 \end{pmatrix} \in M_n(\mathbb{Z})$ and all elements of U_n are $1 \in \mathbb{Z}$.

Consider

$$\mathcal{E}(\Delta) + U_n = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 1 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & 1 \end{pmatrix}.$$

Obviously, $\mathcal{E}(\Delta) + U_n$ is the serial ring with the quiver

$$Q = \left\{ \begin{array}{ccccccc} & 1 & & 2 & & \dots & & n-1 & & n \\ & \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \dots & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet \\ \uparrow & & & & & & & & & \downarrow \\ & \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \dots & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet \end{array} \right\}$$

The quiver Q is the simple cycle $(1\ 2 \dots n)$ and the adjacency matrix $[Q]$ is the permutation matrix which corresponds to the simple cycle $(1\ 2 \dots n)$. Obviously, $(1\ 2 \dots n) = \sigma^{-2}$, where σ is the permutation for Gorenstein matrix $\mathcal{E}(H_n(\mathcal{O}))$. Moreover, the quotient ring H_n/R^2 is the Frobenius ring with Nakayama permutation $\nu = (1\ 2 \dots n)$ (the simple cycle σ^{-1}).

7.2.

Let \mathcal{O} be a discrete valuation ring with the unique maximal ideal $\mathfrak{M} = \pi\mathcal{O} = \mathcal{O}\pi$ and let

$$K_n = K_n(\mathcal{O}) = \begin{pmatrix} \mathcal{O} & \mathfrak{M} & \dots & \mathfrak{M} & \mathfrak{M} \\ \mathfrak{M} & \mathcal{O} & \dots & \mathfrak{M} & \mathfrak{M} \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \mathcal{O} & \mathfrak{M} \\ \mathfrak{M} & \mathfrak{M} & \dots & \mathfrak{M} & \mathcal{O} \end{pmatrix}$$

be a tiled order. Obviously,

$$\mathcal{E}_n = \mathcal{E}(K_n K_n^*) = \mathcal{E}(K_n^* K_n) = \begin{pmatrix} 0 & -1 & \dots & -1 & -1 \\ -1 & 0 & \dots & -1 & -1 \\ \dots & \dots & \dots & \dots & \dots \\ -1 & -1 & \dots & 0 & -1 \\ -1 & -1 & \dots & -1 & 0 \end{pmatrix}$$

and

$$U_n + \mathcal{E}_n = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & \dots & 0 & 1 \end{pmatrix}.$$

Denote $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ (1 is on the i -th place). By lemmas 7.4 and 7.5 irreducible A -lattices Q_i with \mathcal{O} -basis $(\mathcal{O}, \mathcal{O}, \dots, \mathcal{O}, \pi\mathcal{O}, \mathcal{O}, \dots, \mathcal{O})$ ($\pi\mathcal{O}$ is on the i -th place) are relatively injective and has exactly one minimal overmodule $(\mathcal{O}, \mathcal{O}, \dots, \mathcal{O}, \mathcal{O}, \mathcal{O}, \dots, \mathcal{O})$. Let $\sigma : i \rightarrow \sigma(i)$ be a permutation of $\{1, 2, \dots, n\}$. Denote by $I_{n,m} = (\mathfrak{M}^{m_{ij}})$ the two-sided ideal of $K_n(\mathcal{O})$, where $w_{i\sigma(i)} = m + 1$, $w_{ij} = m$, $j \neq \sigma(i)$, $i, j = 1, \dots, n$.

Obviously, $I_{m,n}$ is relatively injective right and left A -lattice and Nakayama permutation of Frobenius quotient ring $K_n(\mathcal{O})/I_{m,n}$ coincides with σ .

Acknowledgements

The first author was partially supported by CNPq and FAPESP of Brazil. The last author was supported by FAPESP of Brazil, and he thanks the Department of Mathematics of the University of São Paulo for its warm hospitality during his visit in 2012.

References

- [1] Kirichenko, V.V. On semiperfect rings of injective dimension one. *Sao Paulo J. Math. Sci.*, v. **1**, no 1, (2007), 111-132.
- [2] Denes, J.; Keedwell, A.D. Latin square and their applications. *Academic Press, New York-London*, 1974, 547 p.
- [3] Hazewinkel, M.; Gubareni, N.; Kirichenko, V. V. Algebras, rings and modules. Vol. 2. *Mathematics and Its Applications (Springer)*, 586. Springer, Dordrecht, 2007. xii+400 p.
- [4] Dokuchaev, M.A.; Kirichenko, V.V.; Zelensky, A.V.; Zhuravlev, V.N. Gorenstein matrices. *Algebra and Discrete Math.* 2005, no. **1**, 8-29.

- [5] Hopkins, C. Rings with minimal condition for left ideals. *Ann. of Mathematics*, v. **40**, 1939, 712-730.
- [6] Levitzki, J. On ring, which satisfy the minimum condition for right-hand ideals. *Compositio Math.*, v. **7**, 1939, 214-222.
- [7] Jategaonkar, V.A. Global dimension of tiled orders over a discrete valuation ring. *Trans. Amer. Math. Soc.*, **196**, 1974, 313-330.
- [8] Tarsy, R.B. Global dimension of orders. *Trans. Amer. Math. Soc.*, vol. **151**, 1970, 335-340.
- [9] Faith, C. Algebra II. Ring Theory. *Springer-Verlag, Berlin-Heidelberg-New York*, 1976.
- [10] Hazewinkel, M.; Gubareni, N. and Kirichenko, V.V. Algebras, Rings and Modules. Vol. **I**, Mathematics and Its Applications, vol. 575. *Kluwer Academic Publishers*, 2004, 380 p.

CONTACT INFORMATION

- | | |
|---|---|
| M. A. Dokuchaev | Departamento de Matemática Univ. de São Paulo, Caixa Postal 66281, São Paulo, SP, 05315-970 – Brazil |
| M. V. Kasyanuk,
V. V. Kirichenko | Faculty of Mechanics and Mathematics, Kiev National Taras Shevchenko Univ., Vladimirska Str., 64, 01033 Kiev, Ukraine |
| M. A. Khibina | In-t of Eng. Thermophysics of NAS of Ukraine, Zheljabova Str., 2A, 03057, Kiev, Ukraine |

Received by the editors: 04.12.2014
and in final form 04.12.2014.