

Morita equivalence of semirings with local units

M. Das, S. Gupta, and S. K. Sardar

Communicated by A. V. Zhuchok

ABSTRACT. In this paper we study some necessary and sufficient conditions for two semirings with local units to be Morita equivalent. Then we obtain some properties which remain invariant under Morita equivalence.

1. Introduction

The classical Morita theory for rings has been recognized as one of the most important and fundamental tools in studying the structure of rings. In 1958 Morita [13] established the Morita equivalence theory for rings with identity. In 1983, Abrams [1] made a first step in extending the theory of Morita equivalence to rings without identity. He considered rings with a commuting set of idempotents such that every element of the ring admits one of these idempotents as a two-sided identity and studied the equivalence of the categories of all unitary left modules of these rings. Ánh and Márki [4] further generalized Abrams' result to rings with local units by weakening the condition of commutativity of idempotents. In the year 2011, Katsov and Nam [10] transferred the ring theoretic approach of Morita equivalence to semirings with identity. In [15], Sardar et al. connected Morita equivalence of semirings with a new and equivalent version of Morita context for semirings. Later Katsov et al. [11], Sardar and Gupta [16, 17] independently studied some properties of semirings

The first author is grateful to CSIR, Govt. of India, for providing research support.
2020 MSC: 16Y60, 16Y99.

Key words and phrases: Morita equivalence, Morita context, Morita invariant, semiring, semimodule.

which remain invariant under Morita equivalence. The aim of this paper is to extend the theory to cover a wider range of semirings namely the semirings with local units in the sense that any two elements of the semiring have a common two sided identity. In order to develop this theory we consider the category R -Sem consisting of all unitary left R -semimodules M i.e., semimodules ${}_R M$ such that $RM = M$, where R is a semiring with local units and say two such semirings R and S to be Morita equivalent if the categories R -Sem and S -Sem are equivalent. Since for a semiring A with identity, A -Sem coincides with the category A -SEM of all left A -semimodules, our notion of Morita equivalence coincides with that of semiring with identity [10]. Consequently, some of the results of Katsov et al. [10] are encompassed in their counterparts obtained here. We organize the paper as follows. In Section 2 we recall some necessary preliminaries on semirings and semimodules. In Section 3 we define locally projective unitary R -semimodule and present some characterizing properties of locally projective generators in semimodule categories. In Section 4 we develop some tools to investigate some necessary and sufficient conditions for R -Sem and S -Sem to be equivalent. Analogously to the case of semirings with identity we show that two semirings with local units R and S are Morita equivalent if and only if there exists a unitary Morita context (R, S, P, Q, τ, μ) with τ, μ surjective. We also identify the semirings with local units which are Morita equivalent to semirings with identity (cf. Prop. 4.14). Finally we conclude the paper by studying some properties of semirings preserved under Morita equivalence in Section 5.

2. Preliminaries

A *semiring*¹ [6] is a nonempty set R on which operations of addition and multiplication have been defined such that (1) $(R, +)$ is a commutative monoid with identity element 0, (2) (R, \cdot) is a semigroup, (3) multiplication distributes over addition from either side, (4) $0r = 0 = r0$ for all $r \in R$. A left R -*semimodule* over a semiring R is a commutative monoid $(M, +, 0_M)$ together with a scalar multiplication from $R \times M$ to M , denoted by $(r, m) \mapsto rm$, which satisfies the following identities: (1) $(r + r')m = rm + r'm$, (2) $r(m + m') = rm + rm'$, (3) $(rr')m = r(r'm)$, (4) $r0_M = 0_M = 0m$ for all $r, r' \in R$ and $m, m' \in M$. Right R -semimodules and R - S -bisemimodules are defined analogously. We will distinguish left and right R -semimodules by writing ${}_R M$ and M_R , respectively. Let M and N

¹Although Golan called it a hemiring, we call it semiring in the present article.

be two left R -semimodules. Then a monoid homomorphism $f : M \rightarrow N$ is called a left R -homomorphism if $f(rm) = rf(m)$ for all $r \in R$ and $m \in M$. The set of all R -morphisms from M to N is denoted by $\text{Hom}_R(M, N)$, in particular $\text{End}_R(M)$ denotes the set of all R -morphisms from M to itself. Right R -homomorphisms and bisemimodule homomorphisms are defined analogously.

A semiring R (semimodule P) is called *additively cancellative* [7] if $a + x = a + y$ implies $x = y$ for all $a, x, y \in R$ (respectively $a, x, y \in P$) and called *additively idempotent* [7] if $a + a = a$ for all $a \in R$ (respectively $a \in P$). If for each element a of a semiring R (semimodule P) there exists an element $b \in R$ (respectively $b \in P$) such that $a + b + a = a$ the semiring (respectively semimodule) is said to be *additively regular* [6]. A semiring R (semimodule P) is said to be *zero-sum free* [7] if $a + b = 0$ implies $a = b = 0$ for all $a, b \in R$ (respectively $a, b \in P$). A nonempty subset I of a semiring R is called an *ideal* [6] of R if $i + j \in I$ and $ri, ir \in I$ for any $i, j \in I$ and $r \in R$. A semiring (semimodule) is said to be *Noetherian* [6] if any ascending chain of ideals (respectively subsemimodules) terminates.

Now we recall some preliminaries related to k -ideals and h -ideals. An ideal I of a semiring R is called a k -ideal [8] (also called subtractive ideal in [6]) of R if for $x \in I, y \in R, x + y \in I$ implies $y \in I$. A subsemimodule N of a semimodule P is called a k -subsemimodule¹ (called subtractive subsemimodule in [6]) of P if for $x \in N, y \in P, x + y \in N$ implies $y \in N$. An ideal I of a semiring R is called an h -ideal [8] of R if for $y_1, y_2 \in I, x, z \in R, x + y_1 + z = y_2 + z$ implies $x \in I$. A subsemimodule N of a semimodule P is called an h -subsemimodule [14] of P if for $y_1, y_2 \in N, x, z \in P, x + y_1 + z = y_2 + z$ implies $x \in N$. The k -closure [8] of an ideal I (a subsemimodule N) is denoted by \bar{I} (respectively \bar{N}) and is defined by $\bar{I} = \{x \in R \mid x + i \in I \text{ for some } i \in I\}$ (respectively $\bar{N} = \{x \in P \mid x + p \in N \text{ for some } p \in N\}$). The h -closure [8] of an ideal I (a subsemimodule N) is denoted by \tilde{I} (respectively \tilde{N}) and is defined by $\tilde{I} = \{x \in R \mid x + y_1 + z = y_2 + z \text{ for some } y_1, y_2 \in I, z \in R\}$ (respectively $\tilde{N} = \{x \in P \mid x + p_1 + z = p_2 + z \text{ for some } p_1, p_2 \in N, z \in P\}$).

An ideal I of a semiring R defines a congruence \mathcal{B}_I on R , called the *Bourne congruence* [6], given by $r\mathcal{B}_I r'$ if and only if there exist $a, a' \in I$ satisfying $r + a = r' + a'$. Similarly I defines another congruence \mathcal{I}_I on R , called the *Iizuka congruence* [6], given by $r\mathcal{I}_I r'$ if and only if there exist $a, a' \in I$ and $s \in R$ satisfying $r + a + s = r' + a' + s$. A congruence ρ on a semiring R is called a *ring congruence* [5] if the factor semiring R/ρ is a

¹In the present article subtractiveness is replaced by k .

ring. A subsemimodule N of a semimodule P defines a congruence \mathcal{B}_N on P , called the *Bourne congruence* [6], given by $p\mathcal{B}_Np'$ if and only if there exist $a, a' \in N$ satisfying $p + a = p' + a'$. Similarly N defines another congruence \mathcal{I}_N on P , called the *Iizuka congruence* [6], given by $p\mathcal{I}_Np'$ if and only if there exist $a, a' \in N$ and $p'' \in P$ satisfying $p + a + p'' = p' + a' + p''$.

For preliminaries on category theory we refer to [2], [9] and [12].

We adopt the following notions from Ánh and Márki [4].

Definition 2.1. Let R be a semiring and $E(R)$ be a set of idempotents of R . Then R is said to be a *semiring with local units* if every finite subset of R is contained in a subsemiring of the form eRe where $e \in E(R)$ or equivalently if for any finite number of elements $r_1, r_2, \dots, r_n \in R$, there exists $e \in E(R)$ such that $er_i = r_i = r_ie$ for all $i = 1, 2, \dots, n$. In this case $E(R)$ is a *set of local units (slu)* of R .

Here we give some examples of semirings with local units.

Example 2.2. 1. Suppose L is a distributive lattice with the least element 0 but with no greatest element¹. Consider L together with the addition $+$ and multiplication \cdot defined by $a + b = \sup\{a, b\}$ and $a \cdot b = \inf\{a, b\}$ respectively, for $a, b \in L$. Then $(L, +, \cdot)$ is a semiring with additive identity 0 but with no multiplicative identity. But it is a semiring with local units, as for any two elements $a, b \in L$, by the absorption law, $a \cdot (a + b) = a = (a + b) \cdot a$ and $b \cdot (a + b) = b = (a + b) \cdot b$, i.e., $a + b$ acts as the common two-sided identity of a and b .

2. Let S be a semiring with identity, X be an infinite set and $R = \{f \mid f : X \rightarrow S \text{ has finite support}\}$. Then R together with the operations $(f + g)(x) := f(x) + g(x)$ and $(fg)(x) := f(x)g(x)$ for $f, g \in R$ and $x \in X$ is a semiring without multiplicative identity. But it is a semiring with local units in view of the following reasons. Suppose $f, g \in R$ with finite supports $\text{supp}(f)$ and $\text{supp}(g)$ respectively, define $h : X \rightarrow S$ by $h(x) = 1$ if $x \in \text{supp}(f) \cup \text{supp}(g)$ and $h(x) = 0$ otherwise, then for $x \in \text{supp}(f)$, $fh(x) = f(x)h(x) = f(x)$ and for $x \in X \setminus \text{supp}(f)$, $fh(x) = f(x)h(x) = 0 \cdot h(x) = 0 = f(x)$. By a similar argument $hf = f$ and hence $fh = f = hf$ and similarly $gh = g = hg$, i.e., h acts as a two-sided identity of f and g .

Definition 2.3. A left R -semimodule M over R is said to be *unitary* if $RM = M$ i.e., for each $m \in M$, there exist $r_1, r_2, \dots, r_n \in R$ and $m_1, m_2, \dots, m_n \in M$ such that $m = r_1m_1 + r_2m_2 + \dots + r_nm_n$.

¹ $(\mathbb{N}, \text{lcm}, \text{gcd})$, $(\mathbb{N}, \text{max}, \text{min})$, where \mathbb{N} is the set of all non-negative integers, are some examples of such lattices.

Remark 2.4. If R is a semiring with slu E and M is a unitary R -semimodule then for each $m \in M$, $m = r_1m_1 + r_2m_2 + \cdots + r_nm_n$ for some $r_1, r_2, \dots, r_n \in R$, $m_1, m_2, \dots, m_n \in M$. Now for $r_1, r_2, \dots, r_n \in R$, there exists $e \in E$ such that $er_i = r_i$ for all $i = 1, 2, \dots, n$, therefore $m = \sum_{i=1}^n r_im_i = \sum_{i=1}^n er_im_i = em$. Thus for every finite subset $M' \subset M$ there exists an $e \in E$ such that $eM' = M'$.

By R -Sem we denote the category of unitary left R -semimodules together with usual R -morphisms.

3. Locally projective generators

In what follows unless otherwise mentioned any semiring is with local units and homomorphisms of semimodules are written opposite the scalars.

Definition 3.1. [6] Let R be a semiring with local units. A semimodule $P \in R$ -Sem is said to be *projective* if for a surjective R -morphism $\phi : M \rightarrow N$ and an R -morphism $\alpha : P \rightarrow N$ in R -Sem there exists an R -morphism $\bar{\alpha} : P \rightarrow M$ satisfying $\bar{\alpha}\phi = \alpha$.

Recall that [10], for any R -semimodule P , the trace ideal $\text{tr}(P) := \sum_{q \in \text{Hom}_R(P, R)} Pq \subseteq R$.

Proposition 3.2. *Let R be a semiring with local units. For any semimodule $P \in R$ -Sem, the following are equivalent:*

- (1) $\text{tr}(P) = R$.
- (2) *There exists a surjective R -morphism $\phi : \bigoplus_I P \rightarrow R$ for some index set I .*
- (3) *For every semimodule $M \in R$ -Sem, there exists a surjective R -morphism $\psi : \bigoplus_\Lambda P \rightarrow M$ for some index set Λ .*

Proof. (1) \Rightarrow (2) Consider the family of all R -morphisms, $f_\alpha : P \rightarrow R$. Now if we set $I = \text{Hom}_R(P, R)$, then the coproduct induced map $f = \bigoplus_I f_\alpha : \bigoplus_I P \rightarrow R$ is a surjective R -morphism since $(\bigoplus_I P)f = \sum_{f_\alpha \in I} Pf_\alpha = \text{tr}(P) = R$.

(2) \Rightarrow (3) Suppose there exists a surjective R -morphism $\phi : \bigoplus_I P \rightarrow R$ for some index set I . Now let $M \in R$ -Sem, then for each $m \in M$ consider the map $\rho_m : R \rightarrow M$ defined by $r \mapsto rm$, then the coproduct induced map $\rho = \bigoplus_{m \in M} \rho_m : \bigoplus_M R \rightarrow M$ is a surjective R -morphism since $(\bigoplus_M R)\rho = \sum_{m \in M} R\rho_m = \sum_{m \in M} Rm = M$. Then the direct sum $\phi' = \bigoplus_M \phi : \bigoplus_M (\bigoplus_I P) \rightarrow \bigoplus_M R$ is a surjection. Hence $\psi = \phi'\rho : \bigoplus_\Lambda P \rightarrow M$ is a surjective R -morphism where $\Lambda = \dot{\bigcup}_M I$ ¹.

¹ $\dot{\bigcup}$ denotes the disjoint union

(2) follows trivially from (3).

(2) \Rightarrow (1) Suppose there exists a surjective R -morphism $\phi : \bigoplus_I P \rightarrow R$ for some index set I . Consider the natural inclusions $\iota_i : P \rightarrow \bigoplus_I P$ for all $i \in I$. Now for each $i \in I$, let $\phi_i = \iota_i \phi$, then $R = (\bigoplus_I P)\phi = \sum_{i \in I} P\phi_i \subseteq \sum_{q \in \text{Hom}_R(P, R)} Pq = \text{tr}(P)$. Hence $\text{tr}(P) = R$. \square

Definition 3.3. A semimodule $P \in R\text{-Sem}$ is said to be a *generator* for the category $R\text{-Sem}$ if P satisfies the equivalent conditions of Prop. 3.2.

Let R be a semiring with local units. Let M be a unitary left R -semimodule and A be a subset of M . Then $RA = \{r_1 a_1 + r_2 a_2 + \cdots + r_n a_n \mid n \in \mathbb{N}, r_i \in R, a_i \in A, \text{ for all } i = 1, 2, \dots, n\}$ is the subsemimodule generated by A . If A generates all of the semimodule M then A is a set of generators for M . A unitary R -semimodule M is said to be *finitely generated* if it has a finite set of generators.

We skip the proof of the following proposition as it is analogous to that of its counterpart in module theory (see [3, Proposition 10.1]).

Proposition 3.4. *If M is a finitely generated unitary left R -semimodule then the following hold:*

- (1) *For every set \mathcal{A} of subsemimodules of M that spans M , there is a finite set $\mathcal{F} \subseteq \mathcal{A}$ that spans M .*
- (2) *Every semimodule that generates M finitely generates M .*

Lemma 3.5. *Retract of a projective unitary R -semimodule is projective.*

Proof. Using the definition of projectivity in $R\text{-Sem}$ (cf. Def. 3.1), the result follows easily from the proof of [9, Prop. 1.7.30] by replacing the notion of epimorphism by surjectivity. \square

The next result is simply a restatement of Prop 17.19 of [6] in the special case of the category $R\text{-Sem}$ of R -semimodules where R is a semiring with local units.

Proposition 3.6. *If $\{P_i \mid i \in \Omega\}$ is a family of unitary left R -semimodules then $P = \bigoplus_{i \in \Omega} P_i$ is projective if and only if each P_i is projective.*

Proposition 3.7. *${}_R P$ is a finitely generated projective unitary semimodule if and only if there exists an idempotent $e \in R$ such that P is a retract of $(Re)^n$, $n \geq 1$.*

Proof. Suppose P is a finitely generated projective unitary semimodule. If $P = \{0\}$ then the zero map $\theta : Re \rightarrow P$ is a retraction in $R\text{-Sem}$. So

we assume that $P \neq \{0\}$ and $\{p_1, p_2, \dots, p_n\}$ is a spanning set of ${}_R P$. Then there exists $e^2 = e \in R$ such that $ep_i = p_i$ for all $i = 1, 2, \dots, n$. Consider $\phi : (Re)^n \rightarrow P$ defined by $(x_1, x_2, \dots, x_n)\phi = \sum_{i=1}^n x_i p_i$. Since for any $p \in P$ there exist $r_1, r_2, \dots, r_n \in R$ such that $p = \sum_{i=1}^n r_i p_i$, $(r_1 e, r_2 e, \dots, r_n e)\phi = \sum_{i=1}^n r_i e p_i = \sum_{i=1}^n r_i p_i = p$. Thus ϕ is onto. Now P being projective there exists $h : P \rightarrow (Re)^n$ such that $h\phi = id_P$. Conversely, suppose $\psi : (Re)^n \rightarrow P$ is a retraction in R -Sem. Let $f : A \rightarrow B$ be a surjection in R -Sem and $g : Re \rightarrow B$ be an R -morphism. Define $\bar{g} : Re \rightarrow A$ by $t \mapsto ta$, where $t \in Re$ and $a \in A$ such that $af = eg$ (if there are more than one $a \in A$ with $af = eg$ then we choose any one of them and fix it throughout). Then $\bar{g}f = g$, hence Re is projective. Therefore by Prop. 3.6, $(Re)^n$ is projective and from Lemma 3.5, ${}_R P$ is projective. Also since $(Re)^n$ has a finite spanning set $\{e_i : i = 1, 2, \dots, n\}$, where each $e_i = (0, \dots, e, \dots, 0)$, with e in the i -th place for all $i = 1, 2, \dots, n$, P is spanned by $\{e_i \psi : i = 1, 2, \dots, n\}$. Thus ${}_R P$ is finitely generated. \square

The notions introduced in the following two definitions are adopted from Ánh and Márki [4].

Definition 3.8. Let I be a partially ordered set such that for each $i, j \in I$ there exists $k \in I$ with $i, j \leq k$ and $(M_i)_{i \in I}$ a family of unitary R -semimodules. Then $(M_i)_{i \in I}$ is said to be a *direct system* if for any $i \leq j$ we have R -morphism $\phi_{ij} : M_i \rightarrow M_j$ such that $\phi_{ii} = 1_{M_i}$ for all $i \in I$ and $\phi_{ij}\phi_{jk} = \phi_{ik}$ for $i \leq j \leq k$.

Moreover a direct system $(M_i)_{i \in I}$ is called a *split direct system* if for each $i \leq j$ in I there exists $\psi_{ji} : M_j \rightarrow M_i$ such that $\phi_{ij}\psi_{ji} = 1_{M_i}$ and $\psi_{kj}\psi_{ji} = \psi_{ki}$ for $i \leq j \leq k$. In this case it follows that $\psi_{ii} = 1_{M_i}$.

Definition 3.9. A unitary R -semimodule M is said to be *locally projective* if it is the direct limit of a split direct system $(M_i)_{i \in I}$ consisting of subsemimodules that are finitely generated projective.

Proposition 3.10. *The R -semimodule ${}_R R$ is a locally projective generator.*

Proof. Let E be a set of local units of R . Define a binary relation \leq on E by $e \leq f$ if and only if $ef = fe = e$. Then clearly \leq is a partial order relation on E and R being a semiring with local units (E, \leq) is an upward directed set. Now for each idempotent $e \in R$ and for each pair $e, f \in R$ with $e \leq f$ consider the map $\psi_{fe} : Rf \rightarrow Re$ given by $r' \mapsto r'e$, where $r' \in Rf$ and the natural inclusion maps $\phi_e : Re \rightarrow R$ and $\phi_{ef} : Re \rightarrow Rf$. Then $(Re)_{e \in E}$ is a split direct system in R -Sem and $R = \varinjlim_E Re$ where Re is finitely

generated projective (as seen in the proof of Prop. 3.7) R -semimodule for each $e \in E$. Hence R is locally projective. Also for any unitary R -semimodule M and for each $m \in M$ consider the map $\rho_m : R \rightarrow M$ defined by $r \mapsto rm$, then we have $\rho = \bigoplus_{m \in M} \rho_m : \bigoplus_M R \rightarrow M$, where $(\bigoplus_M R)\rho = \sum_{m \in M} R\rho_m = \sum_{m \in M} Rm = M$, which implies that ρ is a surjection. Therefore R is a generator in R -Sem. \square

Proposition 3.11. *Let M be a locally projective unitary R -semimodule, then every finitely generated subsemimodule P of M is contained in a finitely generated projective subsemimodule of M .*

Proof. Let M be a locally projective unitary R -semimodule. Then there exists a split direct system (cf. Definition 3.8) $(M_i)_{i \in I}$ of finitely generated projective subsemimodules of M such that $M = \varinjlim M_i$. Let $M' = \dot{\cup} M_i / \rho$, where ρ on $\dot{\cup} M_i$ is given by $(x, i)\rho(y, j)$ if and only if there exists $k \in I$, $i, j \leq k$ such that $x\phi_{ik} = y\phi_{jk}$, where $i, j \in I$, $x \in M_i$, $y \in M_j$. Using the existence of $\psi_{j'i'}$ for each $i', j' \in I$, $i' \leq j'$, it then easily follows that $(x, i)\rho(y, j)$ if and only if $x\phi_{ik} = y\phi_{jk}$ for all $k \in I$, $i, j \leq k$. Now it is a routine matter to verify that M' together with the family of R -morphisms $\phi_i : M_i \rightarrow M'$ given by $x \mapsto [(x, i)]_\rho$ is the direct limit of the split direct system $(M_i)_{i \in I}$. Let P be a subsemimodule of M with a finite spanning set $\{p_1, p_2, \dots, p_n\}$. Then identifying M with M' we have $p_k = [(x_k, i_k)]_\rho$ for each $k = 1, 2, \dots, n$ where $i_k \in I$, $x_k \in M_{i_k}$. Let $t \in I$ such that $i_k \leq t$ for all $k = 1, 2, \dots, n$. Then for each $k = 1, 2, \dots, n$ we have $p_k = x_k\phi_{i_k} = x_k\phi_{i_k t}\phi_t \in M_t\phi_t$. Therefore $P \subseteq M_t\phi_t \cong M_t$, where M_t is a finitely generated projective subsemimodule of M . Hence the proof is complete. \square

We observe that if R and S are semirings with local units and U_S and ${}_R V_S$ are unitary then $\text{Hom}_S(U, V)$ is a left R -semimodule by putting, for $\phi \in \text{Hom}_S(U, V)$ and $r \in R$, $(r\phi)(u) = r\phi(u)$ for $u \in U$. The subsemimodule $R\text{Hom}_S(U, V)$ is the largest unitary R -subsemimodule of $\text{Hom}_S(U, V)$.

Proposition 3.12. *Suppose R is a semiring with slu E . Then $\rho : {}_1 R\text{-Sem} \rightarrow R\text{Hom}_R(R, _)$ is a natural isomorphism where for each $M \in R\text{-Sem}$, $\rho_M : M \rightarrow R\text{Hom}_R(R, M)$ is given by $m \mapsto m\rho_M$ ($r \mapsto rm$). For $M' \in R\text{-Sem}$ and $f \in \text{Hom}_R(M, M')$, $\rho_f : R\text{Hom}_R(R, M) \rightarrow R\text{Hom}_R(R, M')$ is given by $\gamma \mapsto \gamma f$.*

Proof. Clearly ρ_M is an R -morphism. Also the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{\rho_M} & R \operatorname{Hom}_R(R, M) \\ f \downarrow & & \downarrow \rho_f \\ M' & \xrightarrow{\rho_{M'}} & R \operatorname{Hom}_R(R, M') \end{array}$$

since $r((m\rho_M)\rho_f) = r(m\rho_M f) = (rm)f = r(mf) = r((mf)\rho_{M'})$. Hence ρ is a natural transformation. For $M \in R\text{-Sem}$, let $m_1, m_2 \in M$, such that $m_1\rho_M = m_2\rho_M$. Now since there exists $e \in E$ such that $em_1 = m_1, em_2 = m_2$, we have $m_1 = e(m_1\rho_M) = e(m_2\rho_M) = m_2$. Hence ρ_M is injective. Now let $r f \in R \operatorname{Hom}_R(R, M)$ and suppose $(r)f = m \in M$ then for any $t \in R$, $t(m\rho_M) = tm = t((r)f) = (\operatorname{tr})f = t(rf)$ i.e., $m\rho_M = rf$. Thus ρ is a natural isomorphism. \square

Definition 3.13. [10] Let M_R be a right R -semimodule and ${}_R N$ be a left R -semimodule. If F is the free \mathbb{N}_0 -semimodule generated by the cartesian product $M \times N$ and σ is the congruence on F generated by all ordered pairs having the form $((m+m', n), (m, n)+(m', n)), ((m, n+n'), (m, n)+(m, n'))$ and $((mr, n), (m, rn))$ with $m, m' \in M_R, n, n' \in {}_R N$ and $r \in R$, then the factor semimodule F/σ is defined to be the *tensor product* of M and N and is denoted by $M \otimes_R N$. When there is no confusion over the semiring, we denote the tensor product as $M \otimes N$ and the class containing (m, n) by $m \otimes n$.

Proposition 3.14. Suppose R is a semiring with $\operatorname{slu} E$ and $M \in R\text{-Sem}$. Then $R \otimes M \cong M$.

Proof. Suppose R is a semiring with $\operatorname{slu} E$ and M is a unitary R -semimodule. Consider the map $\mu : M \rightarrow R \otimes M$ defined by $m \mapsto e \otimes m$, where $m \in M$ and $e \in E$ such that $em = m$. First we show that the definition is independent of the choice of the idempotent e . Suppose e and f are two idempotents in R such that $em = m = fm$. Let $g \in E$ be a common identity of e and f , then $e \otimes m = ge \otimes m = g \otimes em = g \otimes m$. Similarly $f \otimes m = g \otimes m$, hence $e \otimes m = f \otimes m$. Now it is a routine matter to verify that μ is an R -morphism. Also consider the map $\psi : R \otimes M \rightarrow M$ defined by $r \otimes m \mapsto rm$, where $r \in R$ and $m \in M$. Clearly ψ is a well defined R -morphism. Now for $r \in R, m \in M$, we have

$$(r \otimes m)\psi\mu = (rm)\mu = e \otimes rm = er \otimes m = r \otimes m,$$

where $e \in E$ such that $er = r$, i.e., $erm = rm$. Also

$$m\mu\psi = (g \otimes m)\psi = gm = m,$$

where $g \in E$ such that $gm = m$.

Hence μ is an isomorphism, i.e., $R \otimes M \cong M$. \square

Suppose R is a semiring with $\text{slu } E(R)$ and ${}_R P$ is a unitary semimodule. Let T be a subsemiring of $\text{End}_R P$ having local units $E(T)$ such that $T \text{End}_R P = T$ and $P \in \text{Sem-}T$. Now consider the $T - R$ bisemimodule $Q = T \text{Hom}_R(P, R)R$. Then define:

$$\begin{aligned} \tau : P \otimes Q &\rightarrow R & \text{and} & & \mu : Q \otimes P &\rightarrow T \\ p \otimes q &\mapsto pq & & & q \otimes p &\mapsto qp \quad (p' \mapsto (p'q)p) \end{aligned}$$

It is routine to verify that the maps τ, μ are respectively $R - R$ and $T - T$ bisemimodule morphisms. Also, there is a QPQ -associativity, i.e., for any $q, q' \in Q$ and $p' \in P$, $q(p'q') = (qp')q'$ since for any $p \in P$, $p(q(p'q')) = (pq)(p'q') = ((pq)p')q' = (p(qp'))q' = p((qp')q')$ i.e., $q(pq') = (qp)q'$.

In the notations introduced above, we obtain the following results (cf. Prop. 3.15 - 3.19) characterizing locally projective generators which are the counterparts of Prop. 3.7, 3.10, Theorem 3.11, Prop. 3.12, Corollary 3.13 respectively of [10] in our setting.

Proposition 3.15. *${}_R P$ is locally projective and Pf is finitely generated for all $f \in E(T)$ if and only if $\mu : Q \otimes P \rightarrow T$ is a surjection. Moreover, if μ is a surjection, then it is an isomorphism.*

Proof. For the necessary part, let $f \in E(T)$. Then since Pf is finitely generated, by Prop. 3.11, there exists a finitely generated projective subsemimodule P' of P such that $Pf \subseteq P'$, i.e., $Pf = Pf^2 \subseteq P'f \subseteq Pf$. Therefore $Pf = P'f$, hence it is projective (since $P'f$ being a retract of P' is projective). Therefore by Prop. 3.7, there exists a retraction $\phi : (Re)^n \rightarrow Pf$ for some $n \in \mathbb{N}$, $e^2 = e \in R$ with coretraction $\psi : Pf \rightarrow (Re)^n$, i.e., $\psi\phi = id_{Pf}$. Consider $e_i \in (Re)^n$ with e as the i th coordinate and all others being 0 for each $i = 1, 2, \dots, n$, then for the canonical projections $\pi_i : (Re)^n \rightarrow Re$ we have $\sum_{i=1}^n x\pi_i e_i = x$ for all $x \in (Re)^n$. Let $p_i = e_i\phi$ and $\alpha_i = \pi\psi\pi_i$ for each $i = 1, 2, \dots, n$ where $\pi : {}_R P \rightarrow_R Pf$ is given by $p \mapsto pf$. Now if we put $q_i = f\alpha_i e \in T \text{Hom}_R(P, R)R = Q$, for all $i = 1, 2, \dots, n$. Then for any $p \in P$, we have $pq_i = p(f\alpha_i e) = ((pf)\alpha_i)e =$

$((pf)(\pi\psi\pi_i))e = (pf)(\psi\pi_i)$ for all $i = 1, 2, \dots, n$. Therefore for any $p \in P$,

$$\begin{aligned} p \sum_{i=1}^n q_i p_i &= \sum_{i=1}^n p(q_i p_i) = \sum_{i=1}^n (p q_i) p_i = \sum_{i=1}^n ((pf)(\psi\pi_i))(e_i \phi) \\ &= \left(\sum_{i=1}^n (pf)\psi\pi_i e_i \right) \phi = (pf)\psi\phi = pf, \end{aligned}$$

i.e., $f = \sum_{i=1}^n q_i p_i$. Now for any $t \in T$ there exists an idempotent $f = \sum_{i=1}^n q_i p_i$ such that $t = ft$. Then we have $t = ft = \sum_{i=1}^n q_i p_i t = \mu(\sum_{i=1}^n q_i \otimes p_i t)$. Thus μ is onto. Conversely, for any idempotent $f \in T$, there exist $p_i \in P$, $q_i \in Q$ for $i = 1, 2, \dots, n$ such that $\mu(\sum_{i=1}^n q_i \otimes p_i) = \sum_{i=1}^n q_i p_i = f$. Let $e \in E(R)$ such that $q_i e = q_i$ for all $i = 1, 2, \dots, n$. Then we define $\alpha : (Re)^n \rightarrow Pf$ by $(x_1, x_2, \dots, x_n) \mapsto \sum_{i=1}^n x_i p_i f$ and $\beta : Pf \rightarrow (Re)^n$ by $y \mapsto (y q_1, y q_2, \dots, y q_n)$. Then for $y \in Pf$,

$$\begin{aligned} y\beta\alpha &= (y q_1, y q_2, \dots, y q_n)\alpha = \sum_{i=1}^n (y q_i) p_i f = \sum_{i=1}^n ((y q_i) p_i) f \\ &= \sum_{i=1}^n y(q_i p_i) f = y \left(\sum_{i=1}^n q_i p_i \right) f = y f^2 = y, \end{aligned}$$

i.e., $\beta\alpha = id_{Pf}$. Hence Pf being a retract of $(Re)^n$ is finitely generated projective (by Prop. 3.7). Also, $P = \varinjlim_R Pf$ (can be proved along the same lines as Prop. 3.10). Therefore ${}_R P$ is locally projective.

Now let μ be a surjection and $\mu(\sum_{i=1}^m q_i \otimes p_i) = \mu(\sum_{j=1}^n q'_j \otimes p'_j)$. Since P_T is unitary there exists $f \in E(T)$ such that $p_i f = p_i$, $p'_j f = p'_j$ for all $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. Now by the surjectivity of μ , $f = \sum_{l=1}^k y_l x_l$, where $x_l \in P$, $y_l \in Q$ for all $l = 1, 2, \dots, k$. Then we have

$$\begin{aligned} \sum_{i=1}^m q_i \otimes p_i &= \sum_{i=1}^m q_i \otimes p_i \left(\sum_{l=1}^k y_l x_l \right) = \sum_{i,l} q_i \otimes p_i (y_l x_l) = \sum_{i,l} q_i \otimes (p_i y_l) x_l \\ &= \sum_{i,l} q_i (p_i y_l) \otimes x_l = \sum_{i,l} (q_i p_i) y_l \otimes x_l = \sum_l \left(\sum_i q_i p_i \right) y_l \otimes x_l \\ &= \sum_l \left(\sum_j q'_j p'_j \right) y_l \otimes x_l = \dots = \sum_{j=1}^n q'_j \otimes p'_j, \end{aligned}$$

which proves that μ is injective. Hence μ is an isomorphism. \square

Proposition 3.16. ${}_R P$ is a generator for R -Sem if and only if $\tau : P \otimes Q \rightarrow R$ is a surjection. Moreover, if τ is a surjection, then it is an isomorphism.

Proof. For the necessary part, since ${}_R P$ is a generator, ${}_R R$ is a sum of homomorphic images of P , i.e., every $r \in R$ can be written as $r = \sum_{i=1}^k p_i \phi_i$, $p_i \in P$, $\phi_i \in \text{Hom}_R(P, R)$ for all $i = 1, 2, \dots, k$. Now, since P_T is unitary, there exists $f \in E(T)$ such that $p_i f = p_i$ for all $i = 1, 2, \dots, k$, also there exists $e \in E(R)$ such that $re = r$. Therefore we have

$$\begin{aligned} r &= \left(\sum_{i=1}^k p_i \phi_i \right) e = \sum_{i=1}^k (p_i \phi_i) e = \sum_{i=1}^k p_i (\phi_i e) \\ &= \sum_{i=1}^k (p_i f) (\phi_i e) = \sum_{i=1}^k p_i (f \phi_i e) = \tau \left(\sum_{i=1}^k p_i \otimes f \phi_i e \right), \end{aligned}$$

where $f \phi_i e \in T \text{Hom}_R(P, R) R = Q$. Therefore τ is onto. Conversely, let τ be a surjection then $R = \sum_{q \in Q} P q \subseteq \sum_{q \in \text{Hom}_R(P, R)} P q = \text{tr}(P)$, therefore $R = \text{tr}(P)$. Hence ${}_R P$ is a generator for R -Sem.

Now if we assume τ to be surjective, then the injectivity of τ can be proved in a manner similar to that of μ in Prop. 3.15. \square

Combining the above two results we obtain the following result.

Proposition 3.17. ${}_R P$ is a locally projective generator and ${}_R P f$ is finitely generated for all $f \in E(T)$ if and only if $\mu : Q \otimes P \rightarrow T$ and $\tau : P \otimes Q \rightarrow R$ are T - T and R - R isomorphisms respectively.

Proposition 3.18. Let ${}_R P$ be a locally projective generator for R -Sem and ${}_R P f$ be finitely generated for all $f \in E(T)$. Then the following hold:

- (1) $R \cong (\text{End}_T P) R \cong R \text{End}_T Q$ as semirings.
- (2) $Q := T \text{Hom}_R(P, R) R \cong \text{Hom}_T(P, T) R$ as T - R -bisemimodules.
- (3) $P \cong R \text{Hom}_T(Q, T)$ as R - T -bisemimodules.
- (4) $P \cong (\text{Hom}_R(Q, R)) T$ as R - T -bisemimodules.
- (5) $T \cong (\text{End}_R Q) T$ as semirings.

Proof. (1) Consider the map $\sigma : R \rightarrow \text{End}_T P$ defined by $\sigma(r)(p) := rp$, where $r \in R$, $p \in P$. For any $r_1, r_2 \in R$, $p \in P$, $\sigma(r_1 + r_2)p = (r_1 + r_2)p = r_1 p + r_2 p = \sigma(r_1)(p) + \sigma(r_2)(p) = (\sigma(r_1) + \sigma(r_2))p$, i.e., $\sigma(r_1 + r_2) = \sigma(r_1) + \sigma(r_2)$. Also $\sigma(r_1 r_2)(p) = (r_1 r_2)p = r_1(r_2 p) = \sigma(r_1)\sigma(r_2)(p)$, i.e., $\sigma(r_1 r_2) = \sigma(r_1)\sigma(r_2)$. Thus σ is a semiring morphism. Now let $\sigma(r_1) = \sigma(r_2)$ for some $r_1, r_2 \in R$. Therefore $r_1 p = r_2 p$ for all $p \in P$. Suppose $e \in E(R)$ such that $r_1 = r_1 e$, $r_2 = r_2 e$. Now using Prop. 3.16,

there exist $p_k \in P$, $q_k \in Q$ for $k = 1, 2, \dots, n$ such that $\sum_{k=1}^n p_k q_k = e$. Therefore, $r_1 = r_1 e = r_1 \sum_{k=1}^n p_k q_k = \sum_{k=1}^n (r_1 p_k) q_k = \sum_{k=1}^n (r_2 p_k) q_k = r_2 \sum_{k=1}^n p_k q_k = r_2 e = r_2$. Hence σ is injective. Therefore identifying R with the subsemiring $\sigma(R)$ of $\text{End}_T P$, let $\psi \in (\text{End}_T P)R$, then there exists an idempotent $e' = \sum_{i=1}^m p'_i q'_i \in R$, such that $\psi e' = \psi$. Then for any $p \in P$ we have

$$\begin{aligned} \psi(p) &= (\psi e')p = \psi(e'p) = \psi\left(\sum_{i=1}^m (p'_i q'_i)p\right) = \psi\left(\sum_{i=1}^m p'_i(q'_i p)\right) \\ &= \sum_{i=1}^m \psi(p'_i)(q'_i p) = \sum_{i=1}^m (\psi(p'_i)q'_i)p = \sigma\left(\sum_{i=1}^m (\psi(p'_i)q'_i)\right)(p), \end{aligned}$$

i.e., $\psi = \sigma(\sum_{i=1}^m (\psi(p'_i)q'_i))$. Thus $R \cong (\text{End}_T P)R$ as semirings. Similarly, considering the map $\xi : R \rightarrow \text{End}_T Q$ defined by $\xi(r)(q) := qr$ we can show that $R \cong R \text{End}_T Q$ as semirings.

(2) Define the map $\lambda : Q \rightarrow \text{Hom}_T(P, T)R$ by $\lambda(q)(p) := qp$, where $q \in Q$, $p \in P$. For $q \in Q$ there exists $e' \in E(R)$ such that $qe' = q$, therefore using the QRP -associativity $(\lambda(q)e')p = \lambda(q)(e'p) = q(e'p) = (qe')p = qp = \lambda(q)(p)$, i.e., $\lambda(q) = \lambda(q)e' \in \text{Hom}_T(P, T)R$. That λ is a monoid morphism follows from the fact that μ is a monoid morphism and using the QRP -associativity we have $(t\lambda(q)r)(p) = t\lambda(q)(rp) = t(q(rp)) = t((qr)p) = (tqr)p = \lambda(tqr)(p)$. Thus λ is a T - R morphism. For $q, q' \in Q$, let $\lambda(q) = \lambda(q')$ then for any $p \in P$, $qp = q'p$. Suppose $e^2 = e \in R$ such that $q = qe$, $q' = q'e$. Now, in view of Prop. 3.16, there exist $p_k \in P$, $q_k \in Q$ for $k = 1, 2, \dots, n$ such that $\sum_{k=1}^n p_k q_k = e$. Therefore, $q = qe = q \sum_{k=1}^n p_k q_k = \sum_{k=1}^n (qp_k)q_k = \sum_{k=1}^n (q'p_k)q_k = q' \sum_{k=1}^n p_k q_k = q'e = q'$. Let $\phi \in \text{Hom}_T(P, T)R$, then there exists $e' = \sum_{i=1}^m p'_i q'_i \in E(R)$, such that $\phi e' = \phi$. Then using TQP -associativity, for any $p \in P$ we have

$$\begin{aligned} \phi(p) &= (\phi e')p = \phi(e'p) = \phi\left(\sum_{i=1}^m (p'_i q'_i)p\right) = \phi\left(\sum_{i=1}^m p'_i(q'_i p)\right) \\ &= \sum_{i=1}^m \phi(p'_i)(q'_i p) = \sum_{i=1}^m (\phi(p'_i)q'_i)p = \lambda\left(\sum_{i=1}^m (\phi(p'_i)q'_i)\right)(p), \end{aligned}$$

i.e., $\phi = \lambda(\sum_{i=1}^m (\phi(p'_i)q'_i))$. Thus λ is an isomorphism.

(3),(4) can be proved in a manner similar to (2) and (5) can be proved along the same lines as (1). \square

Proposition 3.19. *Let R_P be a locally projective generator for R -Sem and $R_P f$ be finitely generated for all $f \in E(T)$. Then TQ is T -Sem, P_T is*

$\text{Sem-}T$, $Q_R \in \text{Sem-}R$ are locally projective generators for their respective categories.

Proof. Suppose that ${}_R P \in R\text{-Sem}$ is a locally projective generator for $R\text{-Sem}$ and ${}_R P f$ is finitely generated for all $f^2 = f \in T$. Then by Prop. 3.18, identifying P with $R \text{Hom}_T(Q, T)$ and R with $R \text{End}_T Q$ and using the fact that τ and μ are isomorphisms (Prop. 3.17) and finally applying Prop. 3.17 to ${}_T Q$, we have that ${}_T Q \in T\text{-Sem}$ is a locally projective generator. Similarly P_T , Q_R can be proved to be locally projective generators for their respective categories. \square

4. Morita equivalence and Morita context

Definition 4.1. Let R, S be two semirings with local units. We call R and S to be *Morita equivalent* if the categories $R\text{-Sem}$ and $S\text{-Sem}$ are equivalent, i.e., there exist additive functors $F : R\text{-Sem} \rightarrow S\text{-Sem}$ and $G : S\text{-Sem} \rightarrow R\text{-Sem}$ such that F and G are mutually inverse equivalence functors.

In what follows by equivalence functors we mean additive equivalence functors. In this section we are going to characterize Morita equivalence for semirings with local units (cf. Theorem 4.13). In order to achieve this we first obtain some results below.

Definition 4.2. A unitary bisemimodule ${}_R P_S$ is said to be *faithfully balanced* if the canonical homomorphisms $S \rightarrow \text{End}_R P$ and $R \rightarrow \text{End}_S P$ given by $s \mapsto \rho_s(p \mapsto ps)$ and $r \mapsto \lambda_r(p \mapsto rp)$ respectively, where $s \in S$, $r \in R$, $p \in P$, are injective and identifying R and S with the corresponding subsemirings of endomorphisms of P , $S \text{End}_R P = S$ and $(\text{End}_S P)R = R$.

The following result is analogous to the case of categories of semimodules over a semiring with identity [10] and can be proved in a similar manner.

Lemma 4.3. *Let $F : R\text{-Sem} \rightleftarrows S\text{-Sem} : G$ be an equivalence of the categories $R\text{-Sem}$ and $S\text{-Sem}$, and θ be a surjection in $R\text{-Sem}$. Then $F(\theta)$ is a surjection in $S\text{-Sem}$.*

Lemma 4.4. *Let $F : R\text{-Sem} \rightleftarrows S\text{-Sem} : G$ be an equivalence of the categories $R\text{-Sem}$ and $S\text{-Sem}$, and ${}_R P \in R\text{-Sem}$ be projective. Then $F(P) \in S\text{-Sem}$ is projective, too.*

Proof. By Lemma 4.3, F preserves surjections. So in view of the definition of projectivity in the category of unitary semimodules [cf. Def. 3.1], with relevant modification of the proof of [9, Proposition 5.1.34] by replacing the notion of epimorphism by surjectivity, the result follows easily. \square

Lemma 4.5. *Let $F : R\text{-Sem} \rightleftarrows S\text{-Sem} : G$ be an equivalence of the categories $R\text{-Sem}$ and $S\text{-Sem}$, and ${}_R P \in R\text{-Sem}$ be a generator for $R\text{-Sem}$. Then $F(P) \in S\text{-Sem}$ is a generator for $S\text{-Sem}$.*

Proof. Let $N \in S\text{-Sem}$. Since P is a generator, there exists a surjection $\alpha : P^{(I)} \rightarrow G(N)$ for some non-empty index set I . By Lemma 4.3, $F(\alpha) : F(P^{(I)}) \rightarrow FG(N)$ is a surjection where $FG(N) \cong N$. Also F and G being mutually inverse equivalence functors, by [9, Prop. 5.1.31], G is the right adjoint of F . Then by the dual of [12, Theorem 5.5.1], F preserves direct limits, hence preserves coproducts, i.e., $F(P^{(I)}) \cong F(P)^{(I)}$. Thus N is a homomorphic image of a direct sum of copies of $F(P)$. Hence $F(P)$ is a generator for $S\text{-Sem}$. \square

We skip the proof of Lemma 4.6 and Lemma 4.7 as they can be proved along the same lines as in the case of module theory [3].

Lemma 4.6. *Let $F : R\text{-Sem} \rightarrow S\text{-Sem}$ be a categorical equivalence. Then for each $M, M' \in R\text{-Sem}$ the restriction of F to $\text{Hom}_R(M, M')$, $F : \text{Hom}_R(M, M') \rightarrow \text{Hom}_S(F(M), F(M'))$ is a monoid isomorphism. In particular $F : \text{End}_R(M) \rightarrow \text{End}_S(F(M))$ is a semiring isomorphism.*

Lemma 4.7. *Let $F : R\text{-Sem} \rightarrow S\text{-Sem}$ be an equivalence of the categories $R\text{-Sem}$ and $S\text{-Sem}$, and let ${}_R P \in R\text{-Sem}$ be finitely generated. Then $F(P) \in S\text{-Sem}$ is finitely generated, too.*

Theorem 4.8. *Let $F : R\text{-Sem} \rightleftarrows S\text{-Sem} : G$ be an equivalence of the categories $R\text{-Sem}$ and $S\text{-Sem}$, and let ${}_R P \in R\text{-Sem}$ be a locally projective generator. Then $F(P) \in S\text{-Sem}$ is a locally projective generator, too.*

Proof. By [9, Prop. 5.1.31], G is the right adjoint of F . Then by the dual of [12, Theorem 5.5.1], F preserves direct limits. Using this fact together with Lemmas 4.4, 4.5 and 4.7 we obtain the result. \square

In the following proposition we observe the adjointness of the tensor functor and Hom functor between the categories of unitary semimodules. It is a routine verification so we omit the proof.

Proposition 4.9. *Let R, S be semirings with local units and ${}_S A_R \in S\text{-Sem-}R$, ${}_R B \in R\text{-Sem}$, ${}_S C \in S\text{-Sem}$. Then*

$$\phi : \text{Hom}_S(A \otimes B, C) \rightarrow \text{Hom}_R(B, R\text{Hom}_S(A, C))$$

given by

$$\begin{aligned} \alpha &\mapsto \alpha' : B \rightarrow R\text{Hom}_S(A, C) \\ b &\mapsto b\alpha' : A \rightarrow C \\ &\quad a \mapsto (a \otimes b)\alpha \end{aligned}$$

is a bijective mapping natural in ${}_S A_R$, ${}_R B$, ${}_S C$. In particular, the functor $R\text{Hom}_S(A, -)$ is right adjoint to the functor $A \otimes -$.

The next result is the counterpart of Theorem 4.5 of [10] in this general setting.

Theorem 4.10. *For a functor $F : R\text{-Sem} \rightarrow S\text{-Sem}$ the following statements are equivalent.*

- (1) *F has a right adjoint.*
- (2) *F preserves direct limits.*
- (3) *There exists a unitary S - R -bisemimodule Q such that the functors $Q \otimes - : R\text{-Sem} \rightarrow S\text{-Sem}$ and F are naturally isomorphic.*

Proof. (1) \Rightarrow (2) and (3) \Rightarrow (1) follow from the right analogue of [12, Theorem 5.5.1] and Prop. 4.9 respectively.

(2) \Rightarrow (3) Let $Q := F(R) \in S\text{-Sem}$. Then F induces a right R -semimodule structure on Q with the R -action given by $Q \times R \rightarrow Q$ by $(q, r) \mapsto qF(\rho_r)$ where $\rho_r : R \rightarrow R$ is given by $x \mapsto xr$. In order to show that Q_R is unitary, suppose $q \in Q$. Now $Q = F(\bigcup_{e \in E(R)} Re) = \bigcup_{e \in E(R)} F(Re)$

(since R is a semiring with local units, union coincides in this formula with direct limit and by the hypothesis F preserves direct limits). Therefore $q \in F(Re)$ for some idempotent $e \in R$. Then we have $qe = qF(\rho_e) = q$ (since $\rho_e = 1_{Re}$ implies that $F(\rho_e) = 1_{F(Re)}$). Thus Q is a unitary S - R -bisemimodule. Then the proof follows similarly as in [10, Theorem 4.5]. \square

Theorem 4.11. *Let R and S be Morita equivalent semirings with local units via inverse equivalences $F : R\text{-Sem} \rightarrow S\text{-Sem}$ and $G : S\text{-Sem} \rightarrow R\text{-Sem}$. Set $P = G(S)$ and $Q = F(R)$. Then the following hold:*

- (1) *${}_R P_S, {}_S Q_R$ are unitary faithfully balanced bisemimodules.*
- (2) *${}_R P, P_S, {}_S Q, Q_R$ are locally projective generators.*
- (3) *$F \cong Q \otimes -, G \cong P \otimes -$.*

- (4) $F \cong S \operatorname{Hom}_R(P, -)$, $G \cong R \operatorname{Hom}_S(Q, -)$.
 (5) ${}_R P_S \cong R \operatorname{Hom}_S(Q, S) \cong (\operatorname{Hom}_R(Q, R))S$ and

$${}_S Q_R \cong S \operatorname{Hom}_R(P, R) \cong \operatorname{Hom}_S(P, S)R.$$

Proof. Let $G(S) = P$, then G being an equivalence functor using Lemma 4.6 we have $\operatorname{End}_S S \cong \operatorname{End}_R P$ as semirings. By Prop. 3.12, $S \cong S \operatorname{End}_S S$ as semirings. Since P is a right $\operatorname{End}_R P$ -semimodule, identifying S with the subsemiring $S \operatorname{End}_S S$ of $\operatorname{End}_S S$, P can be considered as a right S -semimodule with the action $P \times S \rightarrow P$ given by $(p, s) \mapsto pG(\rho_s)$ where $\rho_s : S \rightarrow S$ is given by $t \mapsto ts$. That P_S is unitary follows similarly as in the proof of Theorem 4.10. Thus P is a unitary R - S -bisemimodule. Now since S is a locally projective generator, by Theorem 4.8, ${}_R P = G(S)$ is a locally projective generator. In view of Lemma 4.7, $Pf = G(Sf)$ is a finitely generated left S -semimodule for all $f \in E(S)$ and $S \cong S \operatorname{End}_S S \cong S \operatorname{End}_R P$ as semirings. Since ${}_R P$ is a locally projective generator with Pf finitely generated for all $f^2 = f \in S$, using (1) of Prop. 3.18 we have $R \cong (\operatorname{End}_S P)R$ as semirings. Hence ${}_R P_S$ is a faithfully balanced bisemimodule. Similarly $Q = F(R)$ is a unitary faithfully balanced S - R -bisemimodule. Hence (1) is proved.

Since F and G are mutually inverse equivalence functors, they are adjoint to each other [9, Prop. 5.1.31]. Therefore using Theorem 4.10, we obtain $F \cong Q \otimes -$. Similarly $G \cong P \otimes -$. By Prop. 4.9, $Q \otimes -$ is left adjoint to $R \operatorname{Hom}_S(Q, -)$ and $P \otimes -$ is left adjoint to $S \operatorname{Hom}_R(P, -)$. Then by uniqueness of adjoint functors upto natural isomorphism [9, Cor. 5.1.10], we obtain $F \cong Q \otimes - \cong S \operatorname{Hom}_R(P, -)$ and $G \cong P \otimes - \cong R \operatorname{Hom}_S(Q, -)$. This proves (3) and (4).

Now using (4) we obtain, $P = G(S) \cong R \operatorname{Hom}_S(Q, S)$ as R - S -bisemimodule and $Q = F(R) \cong S \operatorname{Hom}_R(P, R)$ as S - R -bisemimodule. Since by (1), Q_R is unitary, using Prop. 3.18 we obtain, $Q = QR \cong S \operatorname{Hom}_R(P, R)R \cong \operatorname{Hom}_S(P, S)R$ as S - R -bisemimodule and also $P \cong (\operatorname{Hom}_R(Q, R))S$ as R - S -bisemimodule, which proves (5). Now (2) clearly follows from Prop. 3.19. \square

Definition 4.12. [15] Let R and S be two semirings and ${}_R P_S$ and ${}_S Q_R$ be an R - S -bisemimodule and an S - R -bisemimodule, respectively and $\tau : P \otimes_S Q \rightarrow R$ and $\mu : Q \otimes_R P \rightarrow S$ be an R - S -bisemimodule homomorphism and an S - R -bisemimodule homomorphism, respectively, such that $\tau(p \otimes q)p' = p\mu(q \otimes p')$ and $\mu(q \otimes p)q' = q\tau(p \otimes q')$ for all $p, p' \in P$ and $q, q' \in Q$. Then the sextuple (R, S, U, V, τ, μ) is called a *Morita context* for semirings.

Moreover, we say that a Morita context is unitary if ${}_R P_S$ and ${}_S Q_R$ are unitary bisemimodules.

Theorem 4.13. *Let R and S be semirings with local units. Then the following are equivalent:*

- (1) R and S are Morita equivalent.
- (2) There exists a faithfully balanced unitary bisemimodule ${}_R P_S$ such that ${}_R P$ is a locally projective generator and ${}_R P f$ is finitely generated for all $f \in E(S)$.
- (3) There exists a unitary Morita context $(R, S, {}_R P_S, {}_S Q_R, \tau, \mu)$ with surjective τ, μ .
- (4) There exists a unitary Morita context $(R, S, {}_R P_S, {}_S Q_R, \tau, \mu)$ with bijective τ, μ .

Proof. (1) \Rightarrow (2) Let $P := G(S)$. Then the proof follows from Theorem 4.11.

(2) \Rightarrow (3) Suppose there exists a unitary bisemimodule ${}_R P_S$ such that ${}_R P$ is a locally projective generator and ${}_R P f$ is finitely generated for all $f \in E(S)$ and $S \cong S \text{End}_R P$ as semirings. Let $Q = S \text{Hom}_R(P, R)R$. Then define:

$$\begin{aligned} \tau : P \otimes Q &\rightarrow R & \text{and} & & \mu : Q \otimes P &\rightarrow S \\ p \otimes q &\mapsto pq & & & q \otimes p &\mapsto qp \quad (p' \mapsto (p'q)p) \end{aligned}$$

It is routine to verify that the maps τ, μ are respectively $R - R$ and $S - S$ morphisms. For any $p' \in P$,

$$\begin{aligned} p' \mu(q \otimes p) q' &= p' ((qp) q') = (p' (qp)) q' = ((p'q)p) q' \\ &= (p'q) (pq') = p' (q(pq')) = p' (q\tau(p \otimes q')), \end{aligned}$$

i.e., $\mu(q \otimes p) q' = q\tau(p \otimes q')$. Also $\tau(p \otimes q) p' = (pq) p' = p(qp') = p\mu(q \otimes p')$.

Consequently $(R, S, {}_R P_S, {}_S Q_R, \tau, \mu)$ is a Morita context. By hypothesis, ${}_R P$ is a locally projective generator and ${}_R P f$ is finitely generated for all $f \in E(S)$ and $S \cong S \text{End}_R P$ as semirings. Hence using Prop. 3.16 and Prop. 3.15, we get that τ, μ are surjections.

(3) \Rightarrow (4) Suppose $(R, S, {}_R P_S, {}_S Q_R, \tau, \mu)$ is a unitary Morita context with surjective τ, μ . Let $\tau(\sum_{i=1}^m p_i \otimes q_i) = \tau(\sum_{j=1}^n p'_j \otimes q'_j)$, where $p_i, p'_j \in P$, $q_i, q'_j \in Q$ for all $i = 1, 2, \dots, m$, $j = 1, 2, \dots, n$. Since Q_R is unitary, there exists an idempotent $e \in R$ such that $q_i e = q_i$, $q'_j e = q'_j$ for all $i = 1, 2, \dots, m$, $j = 1, 2, \dots, n$. Now by the surjectivity of τ , $e =$

$\tau(\sum_{l=1}^k x_l \otimes y_l)$, where $x_l \in P$, $y_l \in Q$ for all $l = 1, 2, \dots, k$. Therefore we have,

$$\begin{aligned} \sum_{i=1}^m p_i \otimes q_i &= \sum_{i=1}^m p_i \otimes q_i \tau\left(\sum_{l=1}^k x_l \otimes y_l\right) = \sum_{i,l} p_i \otimes q_i \tau(x_l \otimes y_l) \\ &= \sum_{i,l} p_i \otimes \mu(q_i \otimes x_l) y_l = \sum_{i,l} p_i \mu(q_i \otimes x_l) \otimes y_l \\ &= \sum_l \sum_i \tau(p_i \otimes q_i) x_l \otimes y_l = \sum_{l=1}^k \tau\left(\sum_{i=1}^m p_i \otimes q_i\right) x_l \otimes y_l \\ &= \sum_{l=1}^k \tau\left(\sum_{j=1}^n p'_j \otimes q'_j\right) x_l \otimes y_l = \dots = \sum_{j=1}^n p'_j \otimes q'_j, \end{aligned}$$

which proves that τ is injective. Similarly μ is also injective.

(4) \Rightarrow (1) Let $(R, S, {}_R P_S, {}_S Q_R, \tau, \mu)$ be a unitary Morita context with bijective τ, μ . Then $P \otimes_S Q \cong R$ and $Q \otimes_R P \cong S$. Therefore for every $M \in R\text{-Sem}$, $P \otimes_S (Q \otimes_R M) \cong (P \otimes_S Q) \otimes_R M \cong R \otimes_R M \cong M$ (cf. Prop. 3.14). Now we consider the class of isomorphisms $\eta = \{\eta_X : P \otimes_S (Q \otimes_R X) \rightarrow_R X \mid X \in R\text{-Sem}\}$. Then η is a natural isomorphism between the identity functor $1_{R\text{-Sem}}$ on the category $R\text{-Sem}$ and the functor $P \otimes_S (Q \otimes_R -)$ as for all ${}_R X, {}_R Y \in R\text{-Sem}$ and $f \in \text{Hom}_R(X, Y)$ the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \uparrow & & \uparrow \\ R \otimes X & \xrightarrow{1_{R \otimes} f} & R \otimes Y \\ \uparrow & & \uparrow \\ P \otimes Q \otimes X & \xrightarrow{1_{P \otimes} 1_Q \otimes f} & P \otimes Q \otimes Y \end{array}$$

Then $P \otimes_S (Q \otimes_R -) \cong 1_{R\text{-Sem}}$. Similarly $Q \otimes_R (P \otimes_S -) \cong 1_{S\text{-Sem}}$. Thus $P \otimes_S - : S\text{-Sem} \rightarrow R\text{-Sem} : Q \otimes_R -$ is an equivalence of the categories $R\text{-Sem}$ and $S\text{-Sem}$. \square

Analogously to Corollary 4.3 of [1], we have the following proposition.

Proposition 4.14. *Let R be a semiring with slu . Then the following are equivalent:*

- (1) R is Morita equivalent to a semiring with identity.
 (2) There exists an idempotent $e \in R$ such that $R = ReR$.

Proof. (1) \Rightarrow (2) Suppose R is Morita equivalent to a semiring S with identity via inverse equivalences $F : R\text{-Sem} \rightleftarrows S\text{-Sem} : G$. Let $P = G(S)$. Since S is a finitely generated projective generator, ${}_R P$ also is a finitely generated projective generator. Now ${}_R P$ being a finitely generated projective unitary R -semimodule, by Prop. 3.7, there exists a surjective R -morphism $\phi : (Re)^m \rightarrow P$ for some idempotent $e \in R$ and $m \in \mathbb{N}$ which implies that Re is a generator for $R\text{-Sem}$. Also since for any $r \in R$, Rr is finitely generated, using Prop. 3.4 there exists a surjective R -morphism $\psi : (Re)^n \rightarrow Rr$ for some $n \in \mathbb{N}$. Therefore there exists $(r_1, r_2, \dots, r_n) \in (Re)^n$ such that $r = (r_1, r_2, \dots, r_n)\psi = r_1e((e, 0, \dots, 0)\psi) + \dots + r_ne((0, \dots, 0, e)\psi) \in ReRr \subseteq ReR$, which is true for any $r \in R$. Therefore $R = ReR$.

(2) \Rightarrow (1) Let $P = Re$. Then clearly P is a finitely generated projective unitary R -semimodule. Also for any $M \in R\text{-Sem}$, for each $m \in M$ consider the map $\rho_m : P \rightarrow M$ defined by $y \mapsto ym$, where $y \in P$, $m \in M$. Then $\rho = \bigoplus_{m \in M} \rho_m : \bigoplus_M P \rightarrow M$, where $(\bigoplus_M P)\rho = \sum_{m \in M} P\rho_m = PM = P(RM) = (PR)M = (ReR)M = RM = M$, which implies that ρ is a surjection. Thus P is a finitely generated projective generator hence a locally projective generator for $R\text{-Sem}$. Now if we take $S = \text{End}_R P = \text{End}_R(Re) = eRe$, then using (2) of Theorem 4.13, R and $S = eRe$ are Morita equivalent semirings. \square

5. Morita invariant properties

In this section we discuss some properties of semirings with local units which remain invariant under Morita equivalence. The results obtained here are nothing but counterparts of the results of [17] in the setting of semirings with local units.

Theorem 5.1. *Let R and S be Morita equivalent semirings with local units via the Morita context $(R, S, {}_R P_S, {}_S Q_R, \tau, \mu)$. Then R is additively cancellative (additively idempotent, additively regular, zero-sum free) if and only if P is additively cancellative (respectively additively idempotent, additively regular, zero-sum free).*

Proof. Let R be additively cancellative and $a, b, c \in P$ such that $a + c = b + c$. Also let $t \in S$ be an idempotent such that $a = at$, $b = bt$. Then the result can be proved in a similar manner to that of [17, Theorem 2.1] by replacing t in place of 1_S . Other parts follow similarly from their corresponding definitions. \square

Theorem 5.2. *Let R and S be Morita equivalent semirings with local units via the Morita context $(R, S, {}_R P_{S,S}, {}_S Q_R, \tau, \mu)$. Then the lattice $\text{Id}(R)$ of ideals of R and the lattice $\text{Sub}(P)$ of subsemimodules of P are isomorphic. Moreover, the isomorphism takes finitely generated ideals to finitely generated subsemimodules and vice-versa.*

Proof. Let us define

$$f : \text{Id}(R) \rightarrow \text{Sub}(P) \quad \text{and} \quad g : \text{Sub}(P) \rightarrow \text{Id}(R)$$

by

$$f(I) := \left\{ \sum_{k=1}^n i_k p_k \mid p_k \in P, i_k \in I \text{ for all } k; n \in \mathbb{N} \right\},$$

and

$$g(N) := \left\{ \sum_{k=1}^n \tau(p_k \otimes q_k) \mid p_k \in N, q_k \in Q \text{ for all } k; n \in \mathbb{N} \right\},$$

respectively. Then with relevant modification of the proof of Theorem 2.2 [17] by replacing the identity by a local unit the rest of the proof can be completed. \square

Remark 5.3. The above result has its counterpart for k -ideals and h -ideals which is analogous to Theorem 2.5 of [17].

Remark 5.4. f and g also preserve k -closure and h -closure.

The following result is an obvious corollary of Theorem 5.2 and the result mentioned in Remark 5.3.

Corollary 5.5. *Let R and S be Morita equivalent semirings with local units via the Morita context $(R, S, {}_R P_{S,S}, {}_S Q_R, \tau, \mu)$. Then R is ideal-simple (k -ideal simple, h -ideal simple) if and only if P is subsemimodule-simple (respectively k -subsemimodule simple, h -subsemimodule simple).*

The following result is the counterpart of Theorem 2.8 of [17] in the present setting.

Theorem 5.6. *Let R and S be Morita equivalent semirings with local units via the Morita context $(R, S, {}_R P_{S,S}, {}_S Q_R, \tau, \mu)$. Then R is Noetherian if and only if P is Noetherian.*

Theorem 5.7. *Let R and S be Morita equivalent semirings with local units via the Morita context $(R, S, {}_R P_{S,S}, Q_R, \tau, \mu)$. Then the lattices $\text{Con}(R)$ and $\text{Con}(P)$ of congruences of R and P respectively are isomorphic. Moreover the isomorphism takes Bourne congruences to Bourne congruences, Iizuka congruences to Iizuka congruences and ring congruences to module congruences and vice-versa.*

Proof. Let us define

$$\alpha : \text{Con}(R) \rightarrow \text{Con}(P) \quad \text{by} \quad \alpha(\rho) := \alpha_\rho^{\text{tr}}$$

and

$$\beta : \text{Con}(P) \rightarrow \text{Con}(R) \quad \text{by} \quad \beta(\sigma) := \beta_\sigma^{\text{tr}},$$

where

$$\alpha_\rho = \left\{ \left(\sum_{k=1}^n r_k p_k, \sum_{k=1}^n r'_k p_k \right) \mid (r_k, r'_k) \in \rho, p_k \in P \text{ for all } k; n \in \mathbb{N} \right\}$$

and

$$\beta_\sigma = \left\{ \left(\sum_{k=1}^n \tau(p_k \otimes q_k), \sum_{k=1}^n \tau(p'_k \otimes q_k) \right) \mid (p_k, p'_k) \in \sigma, q_k \in Q \text{ for all } k; n \in \mathbb{N} \right\}.$$

The rest of the proof is a slight modification of the proof of Theorem 2.10 of [17]. \square

The following result is an obvious corollary of the above theorem.

Corollary 5.8. *Let R and S be Morita equivalent semirings with local units via the Morita context $(R, S, {}_R P_{S,S}, Q_R, \tau, \mu)$. Then R is (Bourne, Iizuka, ring) congruence-simple if and only if P is (Bourne, Iizuka, module) congruence-simple.*

6. Concluding remark

All the above results in section 5 investigate relationship between R and P . But similar relationship can be established between R and Q , S and P , S and Q i.e., Theorems 5.1, 5.2, 5.6, 5.7 have their counterparts for other pairs of the components of Morita equivalent semirings with local units. Since a semiring with identity is also a semiring with local units, Theorems 4.11, 4.13 include some of the results of Theorems 4.6, 4.8 of [15].

Acknowledgement

The authors are very much thankful to the learned referee for his meticulous referring and subsequent detection of some errors both in mathematics as well as presentation whose compliance has made the paper what it is now. The authors are also grateful to Prof. L. Márki and Prof. P. N. Ánh of Alfréd Rényi Institute of Mathematics, Hungary for a necessary discussion in order to resolve one issue raised by the learned referee. Lastly the authors would like to acknowledge the encouragement provided by Prof. Y. Katsov of Hanover College, USA at the time of preparation of the paper.

References

- [1] G. D. Abrams, *Morita equivalence for rings with local units*, Communications in Algebra, *11*(8), 1983, 801-837.
- [2] J. Adamek, H. Herlich and G. Strecker, *Abstract and concrete categories*, John Wiley & Sons, Inc., New York, 1990.
- [3] F. W. Anderson and K. R. Fuller, *Rings and categories of modules*, 2nd ed., Springer, New York, 1992.
- [4] P. N. Ánh and L. Márki, *Morita equivalence for rings without identity*, Tsukuba J. Math., *11*(1), 1987, 1-16.
- [5] S. Ghosh, *A characterization of ring congruences on semirings*, Soochow J. Math., *19*(3), 1993, 305-312.
- [6] J. S. Golan, *Semirings and Their Applications*, Kluwer Academic Publishers, Dordrecht, 1999.
- [7] U. Hebisch and H. J. Weinert, *Semirings : Algebraic Theory and Applications in Computer Science*, World Scientific Publishing, Singapore, 1998.
- [8] U. Hebisch and H. J. Weinert, *Semirings and semifields*, Handbook of Algebra, Vol. 1, 1996, 425-462.
- [9] M. Kilp, U. Knauer and A. V. Mikhaev, *Monoids, acts and categories*, Walter de Gruyter, Berlin, New York, 2000.
- [10] Y. Katsov and T. G. Nam, *Morita equivalence and homological characterization of semirings*, Journal of Algebra and Its Applications, *10*(3), 2011, 445-473.
- [11] Y. Katsov, T. G. Nam and J. Zumbragel, *On simpleness of semirings and complete semirings*, J. Algebra Appl., *13*(6), 2014, 1450015 29pp.
- [12] S. Mac Lane, *Categories for the Working Mathematician*, Springer, New York, 1971.
- [13] K. Morita, *Duality of modules and its applications to the theory of rings with minimum condition*, Science Rep. Tokyo Kyoiku Daigaku Sect., A6, 1958, 85-142.
- [14] T. K. Mukherjee, M. K. Sen and S. Ghosh, *Chain conditions on semirings*, International Journal of Mathematics and Mathematical Sciences, *19*(2), 1996, 321-326.

- [15] S. K. Sardar, S. Gupta and B.C. Saha, *Morita equivalence of semirings and its connection with Nobusawa Γ -semirings with unities*, Algebra Colloquium, 22 (Spec 1), 2015, 985-1000.
- [16] S. K. Sardar and S. Gupta, *Morita invariants of semirings*, J. Algebra Appl., 15(2), 2016, 14pp : 1650023.
- [17] S. K. Sardar and S. Gupta, *Morita invariants of semirings-II*, Asian-European Journal of Mathematics, 11(1), 2018: 1850014.

CONTACT INFORMATION

Monali Das, Department of Mathematics,
Sujit Kumar Jadavpur University, Kolkata, India
Sardar *E-Mail(s):* monali.ju7@gmail.com,
sksardarjumath@gmail.com,
sujitk.sardar@jadavpuruniversity.in

Sugato Gupta Department of Mathematics,
Vidyasagar College for Women, Kolkata,
India
E-Mail(s): sguptaju@gmail.com

Received by the editors: 16.11.2018
and in final form 02.09.2019.