

On the mutation loops of valued quivers

I. Saleh

Communicated by I. V. Protasov

ABSTRACT. A mutation loop of a valued quiver, Q , is a combination of quiver automorphisms and mutations that sends Q to itself. Moreover, it will be called *symmetric* if it sends Q to $\epsilon\sigma(Q)$, $\epsilon \in \{-1, 1\}$ for some permutation σ . A *global mutation loop* of Q is a mutation loop that is symmetric for every quiver in the mutation class of Q . This class of relations contains all the relations of the global mutations group yield from the group action on the mutation class of Q . We identify which quivers have global mutation loops and provide some of them for each case.

1. Introduction

S. Fomin and A. Zelevinsky introduced cluster algebras in [17, 8, 9, 10, 2], and a “geometrical” version was introduced by V. Fock and A. Goncharov in [5, 6]. A cluster algebra is a ring with distinguished sets of generators called “*clusters*”. A cluster comprises two tuples of commutative variables called the \mathcal{A} -variables and \mathcal{X} -variables. Each cluster is paired with a “valued” quiver (skew-symmetrizable matrix) to form what is called a *seed*. The combinatorial operation of producing a seed from an existing one is called *mutation*, which relies mainly on the quiver part of the seed.

I want to thank the anonymous paper reviewer for the helpful suggestions which helped get the paper into the final form. Furthermore, I would like to thank Fang Li and Zongzhu Lin for the very valuable discussions regarding the topic of this paper.

2020 Mathematics Subject Classification: *Primary 13F60, Secondary 05E15.*

Key words and phrases: *cluster algebras, global mutation loops.*

The mutation alters the quiver to produce another one. For a quiver Q , the set of all quivers that can be generated from Q by applying sequences of mutations is called the *mutation class* of Q , and we will denote it by $[Q]$. Mutation classes play vital roles in the cluster algebras theory and its related topics.

The transformation of \mathcal{A} -variables (resp. \mathcal{X} -variables) is called the cluster \mathcal{A} -transformation (resp. cluster \mathcal{X} -transformation). A *mutation loop* of a quiver Q is a mutation sequence, formed of quiver mutations and permutations of vertices, that sends Q to itself, see [12, 7]. A mutation loop defines an autonomous discrete dynamical system as the composition of cluster transformations and permutations of coordinates (equivalently permutations of vertices). The mutation loops form a cluster modular group, which acts on some geometric objects called the cluster \mathcal{A} - and \mathcal{X} -varieties and their tropicalizations by a semifield. The discrete dynamical system induced by a mutation loop occurs in these spaces. For more details, see [12], which provides one of the main motives behind the interest in learning about mutation loops.

The second and more practical reason behind the interest in learning about mutation loops comes from [13], where the action of the *global mutations group* on the classes of mutationally equivalent seeds was studied. Moreover, the authors have proven that cluster automorphisms arise naturally when those classes of “labeled” seeds are considered as orbits under the action of the global mutations group. However, finding the group relations of the global mutation group has yet to receive the attention it deserves despite its importance in understanding the algebraic structure of the group and hence the cluster automorphisms, see [13, 14]. This paper is a step toward figuring out these relations considering the mutations classes as orbits of the group action. The main object of this paper is *global mutations loops*, which are generalized relations on the global mutations group.

A *symmetric mutation loop* μ of Q is a sequence of mutations such that there is a permutation σ where $\mu(Q) = \epsilon\sigma(Q)$, $\epsilon \in \{-1, 1\}$. One can see that such mutations sequence μ is a generalized mutation loop. Notice that, based on the definition of mutation loops as provided in references [12] and [7], one can see that every mutation loop is actually symmetric mutation loop with $\sigma = 1$ and $\epsilon = 1$. The main objective of this article is a particular class of symmetric mutation loops called *global mutation loop*.

Definition 1.1. For a valued quiver Q , a *global mutation loop* is a se-

quence of mutations μ that is a symmetric mutation loop for every quiver $Q' \in [Q]$.

For every single mutation μ_i , we consider the global mutation loop μ_i^2 trivial. The primary purpose of this article is to identify valued quivers that exhibit non-trivial global mutation loops and provide them. These valued quivers belong to the broader class of finite mutation-type quivers. The main results of this article are summarized in Theorem 1.2 below. For more details, see Example 3.2, Lemma 3.3, Proposition 3.7, and Theorem 3.10. To facilitate a better understanding, we will introduce the necessary definitions before presenting the main results.

A quiver Q is called *fully cyclic* if every vertex is in a cycle. The *weight of a quiver Q* is the maximum weight of its edges, and the weight of $[Q]$, denoted by $w([Q])$, is the maximum weight of the quivers in $[Q]$. Since all mutation sequences of a rank two quiver are global mutation loops, we will focus on quivers of rank three or more.

Theorem 1.2. *Let Q be a valued quiver. Then we have the following*

1. *If Q is of rank three, then Q will have global mutation loops only if $w([Q]) = 4$, except the case of*

$$Q = \begin{array}{ccc} & \cdot k & \\ & \uparrow & \searrow (1,3) \\ (2,2) & & \\ & \cdot j & \xleftarrow{(3,1)} \cdot i \end{array}$$

2. *If Q is of rank bigger than or equals to four and $w([Q]) = 4$, then Q has global mutation loops if and only if each quiver $Q' \in [Q]$ is fully cyclic.*

The proofs and arguments are mainly based on Definition 2.2 and Remarks 2.3, which is the valued quiver's version of the mutation of skew-symmetrizable matrices definition. In this version of the definition, mutations are applied at the vertices to alter the valued quiver by changing its edges and weights based on specific rules. Since the paper's primary focus is on quivers and their mutations, we will ignore the frozen vertices of cluster algebras.

The paper is organized as follows: we offer a concise introduction to valued quivers and their mutation. The third section provides an overview of mutation loops, including Example 3.2, Lemma 3.3, Proposition 3.7, and Theorem 3.10, which establish the category of valued

quivers featuring global mutation loops. Additionally, this section includes proofs for the remaining classes that does not have non-trivial global mutation loops.

2. Valued Quivers Mutation

Definitions 2.1. 1. An oriented valued quiver of rank n is a quadruple $Q = (Q_0, Q_1, V, d)$, where

- Q_0 is a set of n vertices labeled by $[1, n]$.
- Q_1 is a set of ordered pairs of vertices, that is $Q_1 \subset Q_0 \times Q_0$ such that; $(i, i) \notin Q_1$ for every $i \in Q_0$, and if $(i, j) \in Q_1$, then $(j, i) \notin Q_1$.
- $V = \{(d_{ij}, d_{ji}) \in \mathbb{N} \times \mathbb{N} \mid (i, j) \in Q_1\}$, V is called the valuation of Q . The weight of an edge $\alpha = (i, j)$ is the product $d_{ij}d_{ji}$ and is denoted by $w_{i,j}$. The weight of Q is given by $w(Q) = \max\{w_{i,j}; (i, j) \in Q_1\}$.
- $d = (d_1, \dots, d_n)$ where d_i is a positive integer for each i , such that $d_i d_{ij} = d_{ji} d_j$ for every $i, j \in [1, n]$.

In the case of $(i, j) \in Q_1$, then there is an arrow oriented from i to j , and in notation, we shall use the symbol $\cdot_i \xrightarrow{(d_{ij}, d_{ji})} \cdot_j$, or $\cdot_i \xrightarrow{w_{i,j}} \cdot_j$ when the emphasis is on the weight of the edge. We will also use $rk(Q)$ for the rank of Q . The quiver Q will be called *simply-laced* if $w(Q) = 1$. We will ignore labeling any edge of weight one. Also, a vertex $i \in Q_0$ is called a *leaf* if there is exactly one vertex j such that $w_{ij} \neq 0$ and $w_{kj} = 0$, for all $k \in [1, n] \setminus \{j\}$.

2. A valued quiver Q' is called *symmetric* to Q if and only if there is a permutation τ such that Q' can be obtained from Q by permuting the vertices of Q using τ such that for every edge $\cdot_i \longrightarrow \cdot_j$ in Q , the valuation (d_{ij}, d_{ji}) is assigned to the edge $\cdot_{\tau(i)} \longrightarrow \cdot_{\tau(j)}$ in Q' .

3. A quiver Q of rank $n > 2$, is said to be a *star quiver* if there is a vertex $i \in Q_0$, such that $w_{ij} \neq 0$, for $j \in [1, n] \setminus \{i\}$ and $w_{kj} = 0$, for every $k \in [1, n] \setminus \{i\}$. In other words, each vertex is a leaf except for exactly one.

We note that every oriented valued quiver corresponds to a skew

symmetrizable matrix $B(Q) = (b_{ij})$ given by

$$b_{ij} = \begin{cases} d_{ij}, & \text{if } (i, j) \in Q_1, \\ 0, & \text{if } i = j, \\ -d_{ij}, & \text{if } (j, i) \in Q_1. \end{cases} \quad (2.1)$$

One can also see that every skew symmetrizable matrix B corresponds to an oriented valued quiver Q such that $B(Q) = B$.

All our valued quivers are oriented, so in the rest of the paper, we will omit the word "oriented". All quivers are of rank n unless stated otherwise. We will also remove the word valued from the term "valued quiver" when there is no confusion.

Definition 2.2 (*Valued quiver mutation*). Let Q be a valued quiver. The mutation $\mu_k(Q)$ at a vertex k is defined through Fomin-Zelevinsky's mutation of the associated skew-symmetrizable matrix. The mutation of a skew symmetrizable matrix $B = (b_{ij})$ on the direction $k \in [1, n]$ is given by $\mu_k(B) = (b'_{ij})$, where

$$b'_{ij} = \begin{cases} -b_{ij}, & \text{if } k \in \{i, j\}, \\ b_{ij} + \text{sign}(b_{ik}) \max(0, b_{ik}b_{kj}), & \text{otherwise.} \end{cases} \quad (2.2)$$

The following remarks provide a set of rules that are adequate to calculate mutations of valued quivers without using their associated skew-symmetrizable matrix.

Remarks 2.3. 1. Let $Q = (Q_0, Q_1, V, d)$ be a valued quiver. The mutation $\mu_k(Q)$ at the vertex k is described using the mutation of $B(Q)$ as follows: Let $\mu_k(Q) = (Q_0, Q'_1, V', d)$, we obtain Q'_1 and V' , by altering Q_1 and V , based on the following rules.

- (a) replace the pairs (i, k) and (k, j) with (k, i) and (j, k) respectively and, in the same manner, switch the components of the ordered pairs of their valuations;
- (b) if $(i, k), (k, j) \in Q_1$, such that neither of (j, i) or (i, j) is in Q_1 (respectively $(i, j) \in Q_1$) add the pair (i, j) to Q'_1 , and give it the valuation $(v_{ik}v_{kj}, v_{ki}v_{jk})$ (respectively change its valuation to $(v_{ij} + v_{ik}v_{kj}, v_{ji} + v_{ki}v_{jk})$);
- (c) if $(i, k), (k, j)$ and (j, i) in Q_1 , then we have three cases

- (i) if $v_{ik}v_{kj} < v_{ij}$, then keep (j, i) and change its valuation to $(v_{ji} - v_{jk}v_{ki}, -v_{ij} + v_{ik}v_{kj})$;
 - (ii) if $v_{ik}v_{kj} > v_{ij}$, then replace (j, i) with (i, j) and change its valuation to $(-v_{ij} + v_{ik}v_{kj}, |v_{ji} - v_{jk}v_{ki}|)$;
 - (iii) if $v_{ik}v_{kj} = v_{ij}$, then remove (j, i) and its valuation.
- (d) d will stay the same in $\mu_k(Q)$.

2. One can see that; $\mu_k^2(Q) = Q$ and $\mu_k(B(Q)) = B(\mu_k(Q))$ at each vertex $k \in [1, n]$ where $\mu_k(B(Q))$ is the mutation of the matrix $B(Q)$. For more information on mutations of skew-symmetrizable matrices, see, for example, [10, 14].

Definitions 2.4. • The *mutation class* of Q is the set of all quivers that can be produced from Q by applying some sequence of mutations, and it will be denoted by $[Q]$.

• A quiver Q is called *fully cyclic* if none of the vertices of Q_0 is a leaf, i.e., each vertex is a vertex in one or more cycles. Furthermore, $[Q]$ is called *fully cyclic* if every quiver in $[Q]$ is fully cyclic.

• Let \mathcal{S}_n be the symmetric group in n letters. One can introduce an action of \mathcal{S}_n in the set of quivers of rank n as follows: for a permutation τ , the quiver $\tau(Q)$ is obtained from Q by permuting the vertices of Q using τ such that for every edge $\cdot_i \longrightarrow \cdot_j$ in Q , the valuation (d_{ij}, d_{ji}) is assigned to the edge $\cdot_{\tau(i)} \longrightarrow \cdot_{\tau(j)}$ in $\tau(Q)$.

• The *weight of the mutation class* $[Q]$ is $w([Q]) = \max\{w(Q'); Q' \in [Q]\}$.

• If $\mu = \mu_{i_1} \dots \mu_{i_k}$ is a sequence of mutations, we will use the notations $\{\mu\} := \{\mu_{i_1}, \dots, \mu_k\}$ and $[\mu_j]_\mu$ for the number of times the single mutation μ_j appears in μ .

Definition 2.5. (**Cluster pattern of $[Q]$**). The *cluster pattern* $\mathbb{T}_n(Q)$ of the mutation class $[Q]$ is a regular n -ary tree whose edges are labeled by the numbers $1, 2, \dots, n$ such that the n edges emanating from each vertex receive different labels. The vertices are assigned to be the elements of $[Q]$ such that the endpoints of any edge are obtained from each other by the quiver mutation in the direction of the edge label.

Definition 2.6. A quiver Q is called *finite mutation type* if the mutation class $[Q]$ contains finitely many quivers.

Definitions 2.7. (Subquivers) [11]. 1. Let Q be a quiver of rank n . We can obtain a sub mutation class of $[Q]$ by fixing a subset I of Q_0 and apply all possible sequences of mutations on the directions of vertices from the set I only. In such case we say Q_I is a subquiver of Q . We will use $Q_I \leq Q$ to say Q_I is a subquiver of Q . The vertices $Q_0 \setminus I$ will be called the *frozen vertices* of Q_I .

2. A subquiver Q_I is said to be of *A-type* if the underlying graph of Q_I is of A_n -type such that $w[Q_I] = 1$.

Examples 2.8. 1. Let

$$Q = \begin{array}{ccccccc} & & \xleftarrow{(2,3)} & & \xrightarrow{(2,3)} & & \xleftarrow{(2,1)} \\ & & & \cdot 3 & & \cdot 2 & & \cdot 7 \\ & & & & \swarrow & \downarrow & & \\ & & & & & & \downarrow & \\ & & & & & & & \cdot 5 \\ & & & \cdot 6 & \xrightarrow{(6,2)} & \cdot 1 & \xrightarrow{(2,3)} & \end{array} \quad (2.3)$$

Consider the subquiver Q_I , with $I = \{1, 2, 3\}$, and $d = (1, 2, 3)$. So, $rk(Q_I) = 3$ and $w(Q_I) = 12$. Here, $(6, 2)$ is the valuation of the edge $\cdot 1 \longrightarrow \cdot 3$. Applying mutation at the vertex 2 produces the following quiver

$$\mu_2(Q_I) = \begin{array}{ccccccc} & & \xleftarrow{(2,3)} & & \xleftarrow{(3,2)} & & \xrightarrow{(1,2)} \\ & & & \cdot 3 & & \cdot 2 & & \cdot 7 \\ & & & & & \uparrow & \swarrow & \\ & & & & & & & \downarrow \\ & & & & & & & \cdot 5 \\ & & & \cdot 6 & \xrightarrow{(2,1)} & \cdot 1 & \xrightarrow{(2,3)} & \end{array}$$

2. Consider the quiver $Q = \cdot n \longleftarrow \cdot n-1 \longleftarrow \cdots \longleftarrow \cdot 2 \xleftarrow{(2,1)} \cdot 1$. An easy exercise is to show that there is a sequence of mutations to move the weight two edge to any other position.

Lemma 2.9. *If Q is simply-laced quiver then for every permutation σ there is a sequence of mutations μ such that $\sigma(Q) = \mu(Q)$, for more details see (Theorem 2.6 in [1]).*

3. Global Mutation loops

Fix a quiver Q with $[Q]$ as its mutation class. Let \mathcal{M} be the set of all reduced sequences (formal words) formed from the elements of the set $\{\mu_1, \dots, \mu_n\}$. Consider the action of the elements of \mathcal{M} on the mutation class $[Q]$ where elements of \mathcal{M} are formed by compositions of mutations.

Definitions and Notations 3.1. A. Let $\mathcal{M}(Q)$ be the group generated by the elements of \mathcal{M} subject to the relations due to its action on $[Q]$. The group $\mathcal{M}(Q)$ is called *the global mutation group* of Q .

B. Consider the following particular types of elements of $\mathcal{M}(Q)$.

1. A sequence of mutations μ is called *homogeneous* if whenever $\mu = \mu^{(1)}\mu^{(2)}$, such that $\mu^{(i)} \neq 1$ for $i = 1, 2$, then we have $\{\mu^{(1)}\} \cap \{\mu^{(2)}\} \neq \emptyset$. Otherwise μ will be called *inhomogeneous* and $\mu^{(i)}$ will be called *isolated subsequence* of mutations of μ for $i = 1, 2$.
2. A sequence of mutations μ will be *full* if $\{\mu\} = \{\mu_1, \dots, \mu_n\}$.
3. Let μ be a mutation sequence such that $\{\mu\}$ contains more than one element. Then μ is called *symmetric mutation loop* of Q if $\mu(Q) = \epsilon\tau(Q)$, $\epsilon \in \{-1, 1\}$ for some permutation τ . The subset of $\mathcal{M}(Q)$ of all symmetric mutation loops is called the *symmetric mutations set* of Q , and it will be denoted by $\mathbf{L}_{Q,S}$.
4. The set of *global symmetric loops* on Q is denoted by $\mathbf{B}_{[Q],S}$ and it will be defined as follows

$$\mathbf{B}_{[Q],S} = \bigcap_{Q' \in [Q]} \mathbf{L}_{Q',S}.$$


We will omit the word symmetric from the term "global symmetric loops" as long as there is no confusion. The rest of the paper aims to identify quivers with non-trivial global loops and provide an explanatory statement for the remaining cases.

Examples 3.2. (Quivers with global loops). In the following, we provide a list of quivers with global loops based on their ranks.

1. Let Q be the quiver $\cdot_i \xrightarrow{(m,n)} \cdot_j$. Then we have $\mathbf{B}_{[Q],S} = \{1, \mu_i\} = \{1, \mu_j\}$.


2. If Q is a periodic quiver, i.e., $[Q] = \{\pm Q\}$, of finite mutation type of rank three, then every mutation sequence is a global loop.

3. Let Q be the quiver



Since the class $[Q]$ is very small, containing only four quivers, it would be a good exercise to show that $\mu_k\mu_i, \mu_i\mu_k, \mu_v\mu_j$ and $\mu_j\mu_v$ are the non-trivial global loops of $[Q]$.

4. Let Q be the quiver




Since the mutation class $[Q] = \{Q\}$, one can see that $\mathbf{B}_{[Q],\sigma}$ is generated by $\{\mu_i, \mu_l, \mu_t, \mu_j, \mu_k\}$.

In the following lemma, we identify all rank three quivers with global loops.

Lemma 3.3. *Let Q be a quiver of rank 3. Then we have the following.*

1. *If $w([Q]) = m$, where $m = 1$, or 2, then $\mathcal{M}(Q)$ does not have any global loops.*
2. *If $w([Q]) = 4$, then $\mathcal{M}(Q)$ contains global loops except the case*



Proof. 1. Assume that $w[Q] = m$; where $m = 1$, or 2. Without loss of generality, let Q be a cyclic quiver with the following underlying graph



One can see that if $\mu \in \mathbf{B}_{Q,S}$ that is not an involution relation, then μ must break and rebuild the cyclic quiver or only reverse of the arrows. So, μ must be one of the mutation sequences in the left-side column in Table 1.1 or formed of one or more of them. Moreover, a short calculation shows that the right-side quivers in Table 1.1 provide counter-examples of quivers where μ does not work as a global relation in $[Q]$. The case of 3 we have $[Q]$ is actually empty assuming it is finite.

Table 1.1: Counter examples of quivers for rank 3 with weight $m = 1$ or 2

μ	Counter examples of quivers
$\mu_i\mu_j\mu_k\mu_i$	$\cdot_i \xleftarrow{\quad} \cdot_k \xrightarrow{m} \cdot_j$
$\mu_i\mu_j\mu_k\mu_i$	$\cdot_i \xrightarrow{\quad} \cdot_k \xrightarrow{m} \cdot_j$
μ_j	$\cdot_i \xleftarrow{m} \cdot_k \xrightarrow{\quad} \cdot_j$

2. Table 1.2 below provides proof of the existence of global loops for all five cases of Q that have global mutation loops. Finally, let

$$Q = \begin{array}{ccc} & \cdot_k & \cdot \\ & \uparrow & \searrow^{(1,3)} \\ (2,2) & & \\ & \cdot_j & \cdot_i \\ & \xleftarrow{(3,1)} & \end{array}$$

One can see that Q is mutationally equivalent to the quiver in (3.1) with $m = 3$. Then from Part 1 above, we have Q has no global mutation loops. \square





The quivers referred to in the following Lemma are denoted as *leading quivers*. The purpose of this lemma is to establish relationships between $w([Q])$ and the structure of certain quivers within $[Q]$.

Lemma 3.4. [15, Lemma 3.11]. *Let Q be a mutationally finite quiver. Then the weight of $[Q]$ is determined as follows*





1. $w[Q] = 2$ if and only if $rk(Q) = 2$ or $[Q]$ contains a quiver that has exactly one edge, say (i, j) , of weight 2, such that the sub-quiver $Q_{\{Q_0 \setminus \{i,j\}\}}$ is of A-type or in otherworld, Q is mutation-equivalent to a quiver of the type B_n, C_n or F_4 .

2. $w[Q] = 3$ if and only if Q is the quiver $\cdot_i \xrightarrow{(3,1)} \cdot_j$.

Table 1.2: Global loops for rank 3 quivers of weight 4

Mutation classes of weight 4	Global mutation loops
	$\mathcal{M}(Q)$
	μ_{ijki}, μ_{ikji}
	μ_{ijki}, μ_{ikji}
	$\mathcal{M}(Q)$


3. If $rk(Q) > 3$ then every quiver $Q' \in [Q]$ of weight 4 satisfies that edges of weight 4 in Q' appear in a cyclic subquiver that is symmetric to one of the following

<p>(a) $Q_{a,x} :$</p> 	<p>(b) $Q_a :$</p> 
<p>(c) $Q_{c,t} :$</p> 	<p>(d) $Q_d :$</p> 


Where $x = 1, 2, 3$ or 4 and $t = 1$ or 2 , such that edges of weight four are not connected outside their cycles.

4. (Quivers with more than one edge of weight 4). If Q' has more than one edge of weight 4, then it is symmetric to one of the following cases.

(a)




(b) $Q_{a,1}^l$:



where the subquiver connecting v and v' is of A -type of rank $l \geq 1$.

(c) Q_a^l or $Q_{a,2}^l$ where the weight 4 is of valuation $(4, 1)$ or $(2, 2)$ respectively with the obvious changes



also the subquiver connecting v and v' is of A -type of rank $l \geq 1$.

Lemma 3.5. Let Q be a quiver with $w([Q]) \leq 2$ and $rk(Q) \geq 3$. Then every global loop of Q is full. In particular, the statement is true if $\mathcal{S}_{n-1} \subset \mathcal{M}(Q)$.

Proof. The proof is divided into two parts based on $w([Q])$. 1. Assume that $w([Q]) = 1$. Then $\mathcal{S}_n \subset \mathcal{M}(Q)$. Let $\mu \in \mathbf{B}_{Q',\mathcal{S}}$ for every $Q' \in [Q]$. Since μ is not an involution, then $\{\mu\}$ contains at least two elements. The case of $\{\mu\}$ containing exactly two elements is obviously can not be a global loop as $rk(Q) \geq 3$, as counter quivers can be found easily. Now

assume that $\{\mu\}$ contains three or more elements. Now, let i, j and k be vertices in $\{\mu\}$. Since $\mathcal{S}_n \subset \mathcal{M}(Q)$, then we can identify a quiver $Q' \in [Q]$ such that $Q'' = \sigma(Q')$, for some permutation σ . Where σ rearranges Q' so that the vertices i, j and k form a subquiver Q^* of Q'' . The subquiver Q^* is isolated from the rest of Q' by a vertex t where $t \notin \{\mu\}$. Here is how Q^* would appear as a subquiver of Q''

$$\cdot i \longrightarrow \cdot j \longrightarrow \cdot k \longrightarrow \cdot t \longrightarrow \cdots . \quad (3.5)$$

One can see that the quiver Q'' is not going to be altered by any mutation sequence that does not contain μ_i, μ_j or μ_k . Therefore, if μ is a global loop, it must contain a sub sequence of mutations that is a global loop on the subquiver Q'' which is impossible, thanks to Part 1 of Lemma 3.3.

2. Assume that $w([Q]) = 2$. Then $[Q]$ contains a quiver Q' with an underlying graph of A_n -type with exactly one single edge of weight 2, thanks to Part 1 of Lemma 3.4. One can see that Q' has the following two features

- (a) The edge of weight two can move to any position on Q' , thanks to Part 2 of Example 2.8;
- (b) Without loss of generality, if we fix the edge of weight two as a leaf, say $\cdot_{n-1} \xrightarrow{2} \cdot_n$. Then the vertices in $[1, n-1]$ are permutable, i.e., $\mathcal{S}_{n-1} \subset \mathcal{M}(Q)$.

Now, if $rk(Q') = 3$, then Q does not have any global loops, thanks to Part 1 of Lemma 3.3. Assume that $rk(Q) \geq 4$. Suppose that μ is not a full global loop. Then at least one vertex, t , that is not represented in $\{\mu\}$. If $t = n$, move the edge of weight 2 to the other end so t becomes free to move anywhere in the A_{n-1} -type subquiver of Q' . So whether t is a leaf or not, we can move it at the position of any other vertex. Therefore, one can isolate a subquiver of three vertices, such as in (3.5), from the rest of the quiver by the vertex t using the fact that $\mathcal{S}_{n-1} \subset \mathcal{M}(Q)$. Then, using the same argument as Part 1 of this lemma, we conclude that such a vertex t does not exist. Then μ must be full. \square

The subsequent lemma is adapted from [15] to better suit the context of this article.

Lemma 3.6. [15, Lemma 3.6]. *A quiver Q is of infinite mutation type if and only if there exists a quiver $Q' \in [Q]$ which contains a 3-cycle $Q^* \leq Q$, where*

$$Q^* = \begin{array}{ccc} \cdot i & & \cdot \\ \uparrow w_2 & \searrow w_1 & \\ \cdot k & \xleftarrow{w_3} & \cdot j \end{array} ,$$

such that $w(\mu_x \mu(Q^)) > w(\mu(Q^*))$ for every sequence of mutations μ and for $x \in \{i, j, k\}$.*

Proposition 3.7. *Let Q be a quiver. Then we have the following.*

1. *If Q is of infinite mutation type, then $\mathcal{M}(Q)$ does not have any non-trivial global loops.*
2. *If Q is a star quiver, then $\mathcal{M}(Q)$ does not have any non-trivial global loop.*

Proof. 1. Suppose that Q is of infinite mutation type. Then $[Q]$ contains a quiver Q' with an unbounded, cyclic subquiver. Hence, from Lemma 3.6, there is a mutation sequence μ such that $w(\mu_i \mu(Q')) > w(\mu(Q))$ for every $i \in [1, n]$. Then $[Q]$ contains a quiver with weights that are impossible to be reproduced using any sequence of mutations. Therefore, $\mathbf{B}_{[Q], \mathcal{S}}$ can not contain any non-involutions loops.

2. The proof of this case is provided for a rank four star quiver, and the higher rank is the same argument. Let Q be a star quiver of rank 4. Then $[Q]$ contains a quiver Q' , which has a subquiver with the following underlying graph



$$\begin{array}{ccc} \cdot i & & \cdot l \\ \downarrow m_3 & \nearrow m_2 & \\ \cdot k & \xleftarrow{m_1} & \cdot j \\ \uparrow & & \\ \cdot t & & \end{array}$$

To finish the proof, Table 1.3 below provides the list of possible mutation loops on Q and a counter example of quivers in $[Q]$ for each case. \square

Lemma 3.8. *If $\mathcal{S}_n \subset \mathcal{M}(Q)$, then every global loop is homogenous, i.e., every global loop does not contain any isolated subsequences.*


Proof. We have two cases: The first is when Q has a subquiver of A_4 -type connected to Q at exactly one vertex. Let $\mu = \mu_{i_1} \cdots \mu_{i_k}$ be a global loop on $[Q]$ with an isolated subsequence μ^* including μ_1, μ_2, μ_3 and μ_4 .

Table 1.3: Counter examples of quivers for rank 4 star quiver


μ	Counter examples
μ_t	
$\mu_t^m (\mu_i \mu_j \mu_l)^k, (\mu_i \mu_j \mu_l)^k \mu_t^m, (\mu_i \mu_j \mu_l)^k$	

Where $m, m_x; x = 1, \dots, 5$, and k are non-negative integers.

Without loss of generality, assume that μ^* appears at the end of μ . Recall that $\mathcal{S}_n \subset \mathcal{M}(Q)$, which makes it possible to use mutations to move and rearrange the vertices and customize the shape of the underlying graph to be in the following shape.




One can see that μ will break and rebuild the triangular cycle. Then the possible choices for the action of μ^* on the subquiver of the vertices $\{i_1, i_2, i_3, i_4\}$ are equivalent to the forms: $\mu_{i_1} \mu_{i_3} \mu_{i_2} \mu_{i_1}$, $\mu_{i_1} \mu_{i_2} \mu_{i_3} \mu_{i_1}$, $\mu_{i_2} \mu_{i_3} \mu_{i_1} \mu_{i_2}$, $\mu_{i_2} \mu_{i_1} \mu_{i_3} \mu_{i_2}$, $\mu_{i_3} \mu_{i_2} \mu_{i_1} \mu_{i_3}$ and $\mu_{i_3} \mu_{i_1} \mu_{i_2} \mu_{i_3}$. Now we will provide a counter-example of a quiver from $[Q]$ for each case. Consider the following quiver from $[Q]$,




If $z = 1$ then $x = 2$ and $y = 3$ for $\mu_{i_1} \mu_{i_3} \mu_{i_2} \mu_{i_1}$ and $x = 3$ and $y = 2$ for $\mu_{i_1} \mu_{i_2} \mu_{i_3} \mu_{i_1}$. And if $z = 2$ then $x = 1$ and $y = 3$ for $\mu_{i_1} \mu_{i_3} \mu_{i_2} \mu_{i_1}$ and $x = 3$ and $y = 1$ for $\mu_{i_1} \mu_{i_2} \mu_{i_3} \mu_{i_1}$. The case of $\mu_{i_3} \mu_{i_2} \mu_{i_1} \mu_{i_3}$ and $\mu_{i_3} \mu_{i_1} \mu_{i_2} \mu_{i_3}$ is similar.

The second case is when Q is of E_6 -type. One can see that the

following quiver, Q' , is in $[Q]$




We use the same argument we used in the first case above, with the quiver Q' , where the counter examples of quivers are encoded in the following quiver.



with x, y , and $z \in \{1, 2, 3\}$, which finishes the proof. □

Remark 3.9. If Q is of finite mutation type quiver, then $[Q]$ is fully cyclic if and only if $[Q]$ contains one of the following quivers



In the following, we will provide the main statement of this article.

Theorem 3.10. *Let Q be a valued quiver. Then we have the following.*

1. *If $w([Q]) \leq 2$ and $rk(Q) \geq 3$, then $[Q]$ has no global loops.*
2. *If $w([Q]) = 3$, then every sequence of mutations is a global loop.*
3. *If $w([Q]) = 4$ and $rk(Q) > 3$, then $\mathcal{M}(Q)$ has global loops if and only if $[Q]$ is fully cyclic. In such case, $[Q]$ contains only two forms of underlying graphs.*

Proof. 1. (a) The case of $rk(Q) = 3$ is proved in Part 1 of Lemma 3.3. The rest of the proof is divided into two cases based on $w([Q])$. Assume that $w([Q]) = 1$. Then $\mathcal{S}_n \subseteq \mathcal{M}(Q)$. Let μ be a global loop of $[Q]$. Then μ is full and homogenous, thanks to Lemma 3.5 and Lemma 3.8. Assume

that $\mu = \mu_2^* \mu_1^*$, where $\{\mu\} = Q_0$ and each of μ_1^* and μ_2^* is a subsequence of mutations of μ . In particular assume that $\{\mu_1^*\} = \{\mu_{i_1}, \mu_{i_2}, \mu_{i_3}, \mu_{i_4}\}$. We have the following remarks on μ

- i. the subsequence μ_2^* could be identity, depending on the rank of Q ;
- ii. in case that $\{\mu_2^*\}$ is not empty, since the subsequence μ_1^* is not isolated then $\{\mu_1^*\} \cap \{\mu_2^*\}$ is not empty;
- iii. if $\{\mu_2^*\}$ is not empty, then there are at least two vertices, say $z \in \{\mu_2^*\}$ and $t \in \{\mu_1^*\}$ such that $[\mu_t]_\mu \neq [\mu_z]_\mu$. To show that, one can use mutations and $S_n \subset \mathcal{M}(Q)$ to customize a quiver $Q' \in [Q]$ with the following features
 - Q' has an underlying graph of A_n -type with two 3-cycles subquivers; one 3-cycle at each end;
 - One of the 3-cycle subquivers contains both z and t as vertices;
 - μ_t appears in μ_2^* before μ_z .

So, μ_t is in the position where applying it breaks its 3-cycle, and it has already been applied within μ_1^* . Furthermore, since μ is redundant-free then $[\mu_t]_\mu > [\mu_z]_\mu$.

Since Q is a simply-laced finite mutation type quiver, then one can find a quiver Q'' in $[Q]$ that has the shape

$$\begin{array}{ccccccc}
 \cdot i_1 & \longleftarrow & \cdot i_2 & & & & \\
 & & \nearrow & & & & \\
 \cdot i_3 & \longleftarrow & \cdot i_4 & \cdots & \longleftarrow & \cdots &
 \end{array} \tag{3.6}$$

There are three possible ways for μ_1^* to can act on the 3-cycle $[t_1, t_2, t_3]$

- Case 1. break and rebuild the 3-cycle completely such that none of the vertices travel outside the 3-cycle over the path of μ , i.e., $\mu(Q'')$ has a subquiver that is symmetric to the subquiver (3.6) with vertices i_1, i_2 , and i_3 .
- Case 2. break and rebuild the 3-cycle completely such that some of the vertices travel outside the 3-cycle and get back over the path of μ , i.e., at least one vertex of the 3-cycle, $[t_1, t_2, t_3]$, is replaced over the path of $\mu(Q'')$.

Case 3. break the 3-cycle and then rebuild it using other vertices, i.e., the 3-cycle appears in $\mu(Q'')$ but in the other end where all vertices i_1, i_2 , and i_3 are replaced.

Case 1: One can see that the 3-cycle (3.6) vertices will not leave it over the path of μ . Then μ will behave as a global loop on the subquiver (3.6). However, simply laced 3-cycles have no global loops, thanks to Part 1 of Lemma 3.3.

Case 2: Suppose that i_2 travels outside the 3-cycle and swaps position with another vertex, say t , and then it returns to the 3-cycle at the end over the path of μ on Q . Fix $i_j \neq i_2$ such that μ_{i_j} to be the last single mutation to appear in μ . Use mutations and the fact that $S_n \subset \mathcal{M}(Q)$ to customize a quiver Q' in $[Q]$ such that it has the same underlying graph of Q with t swapped vertices with $i_j, j \neq 2$, where μ_{i_j} is the last single mutation that appears in μ . Here, the 3-cycle $[t_1, t_2, t_3]$ is formed right after applying μ_{i_j} on the path of $\mu(Q)$. One can see that rebuilding the 3-cycle $[t_1, t_2, t_3]$ in Q' using μ will be interrupted and would not be completed which means $\mu(Q')$ is not symmetric to Q' . Then such a global loop μ does not exist.

Case 3: Without loss of generality assume that μ_2^* is not trivile. In this case, the action of μ causes all the 3-cycles, $[t_1, t_2, t_3]$, vertices to be swapped with other vertices or the 3-cycle that appear in the other end. Without loss of generality, suppose that the new 3-cycle is formed of vertices from $\{\mu_2^*\}$. Since μ is a homogenous mutation loop, then $\{\mu_2^*\} \cap \{\mu_1^*\}$ is non-empty. Now, let μ_t be the last single mutation to appear in μ from $\{\mu_2^*\} \cap \{\mu_1^*\}$. Use mutations and permutations to customize a quiver Q'' in $[Q]$ such that it has the same underlying graph of Q with the vertex t is swapped with z and t_1 becomes a leaf in Q'' . On the path of μ on Q , the 3-cycle will be rebuilt right after applying μ_t and it will stay with no change since it is the last single mutation to appear in $\{\mu_2^*\} \cap \{\mu_1^*\}$. However, on the other hand, the 3-cycle would not be built in $\mu(Q'')$ because over the path of μ on Q'' , the vertex t is exchanged with z and $[\mu_z]_\mu \neq [\mu_t]_\mu$. Hence, $\mu(Q')$ is not symmetric to Q'' .




(b) Assume that $w([Q]) = 2$. Then if there is any global loop, it must be full, thanks to Lemma 3.5. The mutation class of $[Q]$ contains a quiver Q' with an A_n -type underlying graph that has one single edge of weight 2, say $\cdot_i \xrightarrow{2} \cdot_j$. Without loss of generality, assume that the edge of weight two in Q' is a leaf. Using the fact that the weight two

edge can move to any spot and $S_{n-1} \subset \mathcal{M}(Q)$, we can imitate a similar proof of the simply-laced case.

2. Assume that $w([Q]) = 3$. Then Part 2 of Lemma 3.4 guarantees that Q has only two vertices. Therefore, every sequence of mutations of Q is a global loop.

3. The proof of this case is a case-by-case process based on the *leading quivers*, (the quivers in Lemma 3.4). " \Leftarrow ". This case is proved by Table 1.4.

Table 1.4: Global loops for rank 4 quivers of weight 4

Fully cyclic classes of weight 4	Global mutation loops
	$\mu_i, \mu_l, \mu_t, \mu_j, \mu_k$
	μ_{ki}, μ_{jv}
	μ_v, μ_j

" \Rightarrow ". Suppose that Q is not fully cyclic. Then $[Q]$ contains a quiver with at least one leaf vertex. Therefore, Q would be one of the following cases:

$$Q_{c,1}, X_6, X_7, E_6^{(1,1)}, E_7^{(1,1)}, E_8^{(1,1)}, \tilde{Q}_{a,1}^l, \tilde{Q}_{a,2}^l, \tilde{Q}_a^1, \text{ and } \tilde{Q}_a^l, l \geq 2.$$

In the following, we will cover case by case.

- (a) $Q_{c,1}$. If $rk(Q) = 4$, then $[Q]$ contains a star quiver. Then $\mathcal{M}(Q)$ has no global loops, thanks to Part 2 of Proposition 3.7. If $rk(Q) = 5$ then $[Q]$ contains the following quiver

$$Q' = \begin{array}{c} \cdot \\ \cdot \longleftarrow \cdot i_2 \longrightarrow \cdot i_3 \longrightarrow \cdot \\ \uparrow 2 \\ \cdot i_1 \end{array}$$

One can see that if μ is a global loop, it must contain μ_{i_2} . However, the vertex i_2 is a mobile vertex, i.e., one can rearrange the vertices to move i_2 to be a leaf. So, the resulting quiver would not be symmetric to Q' after applying μ .

- (b) $E_6^{(1,1)}, E_7^{(1,1)}, E_8^{(1,1)}, X_6$, and X_7 . One can see that in each of these cases, the mutation class contains a simply-laced quiver. Then they can not have any global loops, thanks to the proof of Part 1 of this theorem and Remark 3.11.
- (c) $\widetilde{Q}_{a,1}^l, l \geq 1$. All of these quivers are mutationally equivalent to simply-laced quivers. Now consider the case of

$$Q = \begin{array}{ccccc} \cdot j' & \xrightarrow{\quad} & \cdot v' & \xleftarrow{\quad} & \cdot v & \xrightarrow{\quad} & \cdot j \\ & \swarrow 4 & \downarrow & & \uparrow & \searrow 4 & \\ & & \cdot k' & & \cdot t_1 & & \end{array}$$

where the weight four edges are either of valuation $(2, 2)$ or $(4, 1)$ with the apparent changes in the weights of other edges. One can see that $[Q]$ contains a simply-laced quiver. Similarly to the previous case, we can mutate Q to a quiver with a simply laced 3-cycle. This process takes us to the same technique used in the previous case.

- (d) $\widetilde{Q}_{a,2}^1, \widetilde{Q}_a^1$. Let Q be the following quiver

$$Q' = \begin{array}{ccccc} \cdot j' & \xrightarrow{2} & \cdot v' & \xrightarrow{\quad} & \cdot v & \xrightarrow{2} & \cdot j \\ & \swarrow 4 & \downarrow 2 & & \uparrow 2 & \searrow 4 & \\ & & \cdot k' & & \cdot k & & \end{array}$$

where the weight 4 is for valuations (2, 2), (1, 4) or (4, 1) with the obvious changes. One can see that $[Q]$ contains the following quivers


$$Q' : \begin{array}{ccccc} \cdot k' & \xrightarrow{2} & \cdot v & \longrightarrow & \cdot v' & \xrightarrow{2} & \cdot j' \\ & & \uparrow 2 & & \uparrow 2 & \searrow 4 & \\ & & \cdot k & & \cdot j & & \end{array}, \text{ and}$$

$$Q'' : \begin{array}{ccccc} \cdot k' & \xrightarrow{2} & \cdot v & \longleftarrow & \cdot v' & \xleftarrow{2} & \cdot j' \\ & & \uparrow 2 & \swarrow 2 & \downarrow 2 & & \\ & & \cdot k & & \cdot j & & \end{array}$$

One can see that applying any sequence of mutations that do not involve μ_v or $\mu_{v'}$ will not alter the edge $\cdot v \longleftarrow \cdot v'$. Then any sequence of mutations that do not contain μ_v or $\mu_{v'}$ would not add or remove edges; hence it would not cause any significant change in the underlying graph of Q . Then, if a sequence of mutations μ does not contain either μ_v or $\mu_{v'}$, it will not be a loop on Q' . Now assume that μ contains μ_v or $\mu_{v'}$, then μ must break and rebuild one or both of the 3-cycles of Q , which would not be a mutation loop of Q'' , considering the positions changes of the vertices of each 3-cycle. Then Q has no global loops.


(e) $\tilde{Q}_{a,2}^l, \tilde{Q}_a^l, l \geq 2$. One can see that the following quiver

$$Q' =$$



is in $[\tilde{Q}_{a,2}^l]$ and $[\tilde{Q}_a^l]$ depending on the valuation of weight 4, where the weight 4 is of valuation either (2, 2) or (4, 1) with the apparent changes in the weights of other edges. Since the subquiver Q^* of Q' , connecting the vertices v and v' is simply laced, then any global loop containing one of the vertices of Q^* has to contain all the

vertices of Q_0^* . Where Q_0^* is the following 3-cyclic subquiver.



One can see that any global loop must alter the 3-cycle, Q_0^* . This argument guarantees that any global loop would contain a subsequence that breaks and rebuilds the subquiver Q_0^* . However, any simply laced quiver of three vertices has no global loop. Therefore, such a global loop μ does not exist.

□


Remarks 3.11. 1. If $[Q]$ contains a simply-laced quiver then $[Q]$ does not have global loops.

2. The case of $rk(Q) = 3$ and $w([Q]) = 4$ is covered in Lemma 3.3.


Proof. The proof of both parts are corollaries of the proof of Part 1 of Theorem 3.10. □

Some exceptional finite mutation types $E_6^{(1,1)}, E_7^{(1)}, E_8^{(1)}$.


$E_6^{(1,1)}$:



$E_7^{(1,1)}$:



$E_8^{(1,1)}$:



References

- [1] A. Alkhezi and I. Saleh, *On the Pentagon Relations of Valued Quivers*, International Journal of Algebra, Vol. 9, no. 8, 2015, pp. 355-369.
- [2] A. Berenstein, S. Fomin and A. Zelevinsky, *Cluster Algebras III: Upper Bounds and Double Bruhat Cells*, Duke Math J., Vol. 126, no. 1, 2005, pp. 1-52.
- [3] H. Derksen, T. Owen, *New graphs of finite mutation type*, Electron. J. Combin. 15 (2008), R139.
- [4] A. Felikson, M. Shapiro, P. Tumarkin, *Cluster algebras of finite mutation type via unfoldings*, Int. Math. Res. Notices Vol. 2012, Issue 8, 2012, pp. 1768–1804.
- [5] V.V. Fock, A.B. Goncharov, *Cluster ensembles, quantization and the dilogarithm*, Ann. Sci. Éc. Norm. Supér. 42(6), 2009, pp. 865–930.
- [6] V.V. Fock, A.B. Goncharov, *Cluster ensembles, quantization and the dilogarithm. II. The intertwiner*, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. 1, pp. 655–673; Progr. Math., 269, Birkhäuser Boston, Inc., Boston (2009).
- [7] V.V. Fock, A. Marshakov, *Loop groups, clusters, dimers and integrable systems*, Geometry and quantization of moduli spaces, pp. 1–66; Adv. Courses Math. CRM Barcelona, Birkhauser/Springer, Cham (2016).
- [8] S. Fomin, A. Zelevinsky, *Cluster algebras I: Foundations*, J. Amer. Math. Soc. 15, 2002, pp. 497-529.
- [9] S. Fomin, A. Zelevinsky, *Cluster algebras II: Finite type classification*, Inven. Math. 154, 2003, pp. 63-121.
- [10] S. Fomin, A. Zelevinsky, *Cluster algebras IV: Coefficients*, Compositio Mathematica, 143(1), 2007, pp. 112-164.
- [11] M. Huang, F. Li and Y. Yang, *On structure of cluster algebras of geometric type I: In view of sub-Seeds and seed homomorphisms*, Science China Mathematics, 61, Issue 5, pp. 831-854.
- [12] T. Ishibashi, S. Kano, *Algebraic entropy of sign-stable mutation loops*. Geom Dedicata, 214, 2021, pp. 79-118.
- [13] A. King and M. Pressland, *Labelled seeds and the mutation group*, Math. Proc. Cambridge Philos. Soc. 163 (2017), no. 2, 193-217. Issue 5, pp. 831-854.
- [14] I. Saleh, *Exchange maps of cluster algebras*, International Electronic J. of Algebra 16, 2014, pp. 1-15.

- [15] I. Saleh, *Symmetric mutations algebras in the context of sub-cluster algebras*, arXiv:2105.08211, to appear in *Journal of Algebra and its Application*, Vol. 23 No. 8 (August 2024).
- [16] A. Seven, *Reconsizing cluster algebras of finite type*, *Electron. J. Combin* 14 (2007), # R3 35pp.
- [17] A. Zelevinsky. *Cluster algebras: Notes for 2004 IMCC* (Chonju, Korea, August (2004), arXiv:math.RT/0407414.

CONTACT INFORMATION

Ibrahim Saleh

The University of Wisconsin Whitewater,
800 W Main St, Whitewater, WI 53190,
USA

E-Mail: salehi@uww.edu

URL:

Received by the editors: 03.04.2023
and in final form 26.09.2023.