

On the semigroup of monoid endomorphisms of the semigroup $\mathcal{C}_+(a, b)$

Oleg Gutik and Sher-Ali Penza

Communicated by V. Mazorchuk

*Dedicated to Professor Yu. A. Drozd
on the occasion of his 80th birthday*

ABSTRACT. Let $\mathcal{C}_+(a, b)$ be the submonoid of the bicyclic monoid which is studied in [8]. We describe monoid endomorphisms of the semigroup $\mathcal{C}_+(a, b)$ which are generated by the family of all congruences of the bicyclic monoid and all injective monoid endomorphisms of $\mathcal{C}_+(a, b)$.

Introduction

We shall follow the terminology of [1, 2, 7]. By ω we denote the set of all non-negative integers, by \mathbb{N} the set of all positive integers.

A semigroup S is called *inverse* if for any element $x \in S$ there exists a unique $x^{-1} \in S$ such that $xx^{-1}x = x$ and $x^{-1}xx^{-1} = x^{-1}$. The element x^{-1} is called the *inverse of $x \in S$* . If S is an inverse semigroup, then the function $\text{inv}: S \rightarrow S$ which assigns to every element x of S its inverse element x^{-1} is called the *inversion*.

If S is a semigroup, then we shall denote the subset of all idempotents in S by $E(S)$. If S is an inverse semigroup, then $E(S)$ is closed under multiplication and we shall refer to $E(S)$ as a *band* (or the *band of S*). Then the semigroup operation on S determines the following partial order

2020 Mathematics Subject Classification: 20M15, 20M20.

Key words and phrases: *endomorphism, injective, bicyclic semigroup, sub-semigroup, direct product, semidirect product.*

\preceq on $E(S)$: $e \preceq f$ if and only if $ef = fe = e$. This order is called the *natural partial order* on $E(S)$. A *semilattice* is a commutative semigroup of idempotents.

If S is an inverse semigroup then the semigroup operation on S determines the following partial order \preceq on S : $s \preceq t$ if and only if there exists $e \in E(S)$ such that $s = te$. This order is called the *natural partial order* on S [9].

The bicyclic monoid $\mathcal{C}(p, q)$ is the semigroup with the identity 1 generated by two elements p and q subjected only to the condition $pq = 1$. The semigroup operation on $\mathcal{C}(p, q)$ is determined as follows:

$$q^k p^l \cdot q^m p^n = q^{k+m-\min\{l,m\}} p^{l+n-\min\{l,m\}}.$$

It is well known that the bicyclic monoid $\mathcal{C}(p, q)$ is a bisimple (and hence simple) combinatorial E -unitary inverse semigroup and every non-trivial congruence on $\mathcal{C}(p, q)$ is a group congruence [1].

Let $\mathfrak{h}: S \rightarrow T$ be a homomorphism of semigroups. Then for any $s \in S$ and $A \subseteq S$ by $(s)\mathfrak{h}$ and $(A)\mathfrak{h}$ we denote the images of s and A , respectively, under the homomorphism \mathfrak{h} . Also, for any $t \in T$ by $(s)\mathfrak{h}^{-1}$ we denote the full preimage of s under the map \mathfrak{h} . A homomorphism $\mathfrak{h}: S \rightarrow T$ of monoids which preserves the unit elements of S is called a *monoid homomorphism*. A homomorphism $\mathfrak{h}: S \rightarrow S$ of a semigroup (a monoid) is called an endomorphism (a monoid endomorphism) of S , and in the case when \mathfrak{h} is an isomorphism then \mathfrak{h} is said to be an *automorphism* of S .

It is well-known that every automorphism of the bicyclic monoid $\mathcal{C}(p, q)$ is the identity self-map of $\mathcal{C}(p, q)$ [1], and hence the group $\mathbf{Aut}(\mathcal{C}(p, q))$ of automorphisms of $\mathcal{C}(p, q)$ is trivial. In [6] all endomorphisms of the bicyclic semigroup are described and it is proved that the semigroups $\mathbf{End}(\mathcal{C}(p, q))$ of all endomorphisms of the bicyclic semigroup $\mathcal{C}(p, q)$ is isomorphic to the semidirect products $(\omega, +) \rtimes_{\varphi} (\omega, *)$, where $+$ and $*$ are the usual addition and the usual multiplication on ω .

Subsemigroups of then bicyclic monoid were studied in [3,4,8]. In [8] the following anti-isomorphic subsemigroups of the bicyclic monoid

$$\mathcal{C}_+(a, b) = \{b^i a^j \in \mathcal{C}(a, b) : i \leq j, i, j \in \omega\}$$

and

$$\mathcal{C}_-(a, b) = \{b^i a^j \in \mathcal{C}(a, b) : i \geq j, i, j \in \omega\}$$

are studied. In the paper [5] topologizations of the semigroups $\mathcal{C}_+(a, b)$ and $\mathcal{C}_-(a, b)$ are studied.

Later in this paper by $\mathfrak{End}(\mathcal{C}_+(a, b))$ we denote the semigroup of all monoid endomorphisms of the semigroup $\mathcal{C}_+(a, b)$.

In this paper we describe monoid endomorphisms of the semigroup $\mathcal{C}_+(a, b)$ which are generated by the family of all congruences of the bicyclic monoid and all injective monoid endomorphisms of $\mathcal{C}_+(a, b)$.

1. On monoid endomorphisms of $\mathcal{C}_+(a, b)$ which are restrictions of homomorphisms of the bicyclic monoid

In [6] was proved that every monoid endomorphism $\varepsilon: \mathcal{C}(a, b) \rightarrow \mathcal{C}(a, b)$ of the bicyclic monoid is one of the following forms:

- (i) $\varepsilon = \lambda_k$ for some positive integer k , where $(b^i a^j) \lambda_k = b^{ki} a^{kj}$ for any $i, j \in \omega$;
- (ii) $\varepsilon = \lambda_0$ is the annihilating endomorphism of $\mathcal{C}(a, b)$, i.e., $(b^i a^j) \lambda_0 = 1$ for any $i, j \in \omega$.

Simple verifications show that in the both cases each of these monoid endomorphisms of the bicyclic semigroup induces the monoid endomorphism of $\mathcal{C}_+(a, b)$, which we denote by the similar way:

$$\lambda_k: \mathcal{C}_+(a, b) \rightarrow \mathcal{C}_+(a, b), b^i a^j \mapsto b^{ki} a^{kj}, i, j \in \omega$$

for some $k \in \omega$.

For any $k_1, k_2 \in \omega$ we have that

$$(b^i a^j) (\lambda_{k_1} \circ \lambda_{k_2}) = ((b^i a^j) \lambda_{k_1}) \lambda_{k_2} = (b^{k_1 i} a^{k_1 j}) \lambda_{k_2} = b^{k_1 k_2 i} a^{k_1 k_2 j}, i, j \in \omega.$$

This implies that $\lambda_{k_1} \circ \lambda_{k_2} = \lambda_{k_1 k_2}$ for all $k_1, k_2 \in \omega$, and hence the set $\{\lambda_k: k \in \omega\}$ of endomorphisms of $\mathcal{C}_+(a, b)$ is closed under the operation of composition.

By $\mathfrak{End}_{\langle \lambda \rangle}(\mathcal{C}_+(a, b))$ we denote the subsemigroup of $\mathfrak{End}(\mathcal{C}_+(a, b))$, which is generated by the family $\{\lambda_k: k \in \omega\}$ of endomorphisms of the monoid $\mathcal{C}_+(a, b)$.

Proposition 1. *The semigroup $\mathfrak{End}_{\langle \lambda \rangle}(\mathcal{C}_+(a, b))$ is isomorphic to the multiplicative semigroup $(\omega, *)$ of non-negative integers.*

Proof. We define the map $\mathfrak{J}: \mathfrak{End}_{\langle \lambda \rangle}(\mathcal{C}_+(a, b)) \rightarrow (\omega, *)$ by the formula $(\lambda_k) \mathfrak{J} = k$ for any $\lambda_k \in \mathfrak{End}_{\langle \lambda \rangle}(\mathcal{C}_+(a, b))$. The above arguments and simple verifications show that so defined map \mathfrak{J} is a semigroup isomorphism. \square

It is well known that any inverse semigroup S admits the *smallest* (minimal) group congruence \mathfrak{C}_{mg} : $a\mathfrak{C}_{\text{mg}}b$ if and only if there exists $e \in E(S)$ such that $ea = eb$ (see [7]). The smallest group congruence \mathfrak{C}_{mg} on the bicyclic semigroup $\mathcal{C}(a, b)$ is determined in the following way: $b^{i_1}a^{j_1}\mathfrak{C}_{\text{mg}}b^{i_2}a^{j_2}$ if and only if $i_1 - j_1 = i_2 - j_2$ [7]. Since the quotient semigroup $\mathcal{C}(a, b)/\mathfrak{C}_{\text{mg}}$ is isomorphic to the additive group of integers $\mathbb{Z}(+)$, the natural homomorphism $\mathfrak{h}_{\mathfrak{C}_{\text{mg}}}: \mathcal{C}(a, b) \rightarrow \mathcal{C}(a, b)/\mathfrak{C}_{\text{mg}}$ generates the homomorphism $\mathfrak{h}_{\mathfrak{C}_{\text{mg}}}: \mathcal{C}(a, b) \rightarrow \mathbb{Z}(+)$ by the formula $(b^i a^j)\mathfrak{h}_{\mathfrak{C}_{\text{mg}}} = j - i$, $i, j \in \omega$. By $(\omega, +)$ we denote the additive group of non-negative integers. This implies that the restriction $\mathfrak{h}_{\mathfrak{C}_{\text{mg}}}|_{\mathcal{C}_+(a, b)}: \mathcal{C}_+(a, b) \rightarrow (\omega, +)$ of the homomorphism $\mathfrak{h}_{\mathfrak{C}_{\text{mg}}}$ is a homomorphism, as well.

Lemma 1. For any $i, j, k \in \omega$ with $j \geq i$ the set

$$S_{i,j,k} = \{b^i a^i\} \cup \{(b^j a^{j+k})^n : n \in \mathbb{N}\}$$

with the induced semigroup operation from the bicyclic monoid $\mathcal{C}(a, b)$ is isomorphic to the semigroup $(\omega, +)$.

Proof. We define the mapping $\mathfrak{J}_{i,j,k}: (\omega, +) \rightarrow S_{i,j,k}$ by the formula

$$(n)\mathfrak{J}_{i,j,k} = \begin{cases} b^i a^i, & \text{if } n = 0; \\ (b^j a^{j+k})^n, & \text{if } n > 0. \end{cases}$$

Simple verifications show that such defined map $\mathfrak{J}_{i,j,k}$ is a bijective homomorphism. □

Definition 1. For arbitrary $l \in \omega$ and $m \in \mathbb{N}$ we define the map $\sigma_{l,m}: \mathcal{C}_+(a, b) \rightarrow S_{0,l,m}$ by the formula $(b^i a^j)\sigma_{l,m} = ((b^i a^j)\mathfrak{h}_{\mathfrak{C}_{\text{mg}}})\mathfrak{J}_{0,l,m}$, $i \leq j$, $i, j \in \omega$. Since $\mathfrak{h}_{\mathfrak{C}_{\text{mg}}}$ and $\mathfrak{J}_{0,l,m}: (\omega, +) \rightarrow S_{0,l,m}$ are homomorphisms, $\sigma_{l,m}$ is a homomorphism, too. Simple verifications show that

$$(b^i a^j)\sigma_{l,m} = \begin{cases} 1, & \text{if } i = j; \\ b^l a^{l+m(j-i)}, & \text{if } i < j \end{cases}$$

for all $i, j \in \omega$.

We observe that every elements of the semigroup $\mathcal{C}_+(a, b)$ can be represented in the form $b^i a^{i+j}$ for some $i, j \in \omega$. Then for any positive integer n we have that

$$\begin{aligned} (b^i a^{i+j})^n &= b^i a^{i+j} \cdot b^i a^{i+j} \cdot (b^i a^{i+j})^{n-2} = \\ &= b^i a^{i+2j} \cdot (b^i a^j)^{n-2} = \\ &= \dots = \\ &= b^i a^{i+nj}. \end{aligned}$$

Lemma 2. $\sigma_{l_1, m_1} \circ \sigma_{l_2, m_2} = \sigma_{l_2, m_1 m_2}$ for arbitrary $l_1, l_2 \in \omega$ and $m_1, m_2 \in \mathbb{N}$.

Proof. Fix an arbitrary $b^i a^j \in \mathcal{C}_+(a, b)$, $i, j \in \omega$. Then we have that

$$\begin{aligned} ((b^i a^j) \sigma_{l_1, m_1}) \sigma_{l_2, m_2} &= \begin{cases} (1) \sigma_{l_2, m_2}, & \text{if } i = j; \\ ((b^{l_1} a^{l_1 + m_1})^{j-i}) \sigma_{l_2, m_2}, & \text{if } i < j \end{cases} = \\ &= \begin{cases} 1, & \text{if } i = j; \\ (b^{l_1} a^{l_1 + (j-i)m_1}) \sigma_{l_2, m_2}, & \text{if } i < j \end{cases} = \\ &= \begin{cases} 1, & \text{if } i = j; \\ (b^{l_2} a^{l_2 + m_2})^{(j-i)m_1}, & \text{if } i < j \end{cases} = \\ &= \begin{cases} 1, & \text{if } i = j; \\ b^{l_2} a^{l_2 + (j-i)m_1 m_2}, & \text{if } i < j \end{cases} = \\ &= (b^i a^j) \sigma_{l_2, m_1 m_2}. \end{aligned}$$

□

By $\mathbf{End}_{\langle \sigma \rangle}(\mathcal{C}_+(a, b))$ we denote the subsemigroup of $\mathbf{End}(\mathcal{C}_+(a, b))$, which is generated by the family $\{\sigma_{l, m} : l, m \in \omega, m > 0\}$ of endomorphisms of the monoid $\mathcal{C}_+(a, b)$.

By $\mathfrak{RZ}(\omega)$ we denote the set ω with the right-zero multiplication, i.e., $xy = y$ for all $x, y \in \omega$, and by $(\mathbb{N}, *)$ the multiplicative semigroup of positive integers. We define the map $\mathfrak{J} : \mathbf{End}_{\langle \sigma \rangle}(\mathcal{C}_+(a, b)) \rightarrow \mathfrak{RZ}(\omega) \times (\mathbb{N}, *)$ by the formula $(\sigma_{l, m})\mathfrak{J} = (l, m)$, $l \in \omega, m \in \mathbb{N}$. Lemma 2 implies that such defined map \mathfrak{J} is a semigroup homomorphism, and moreover \mathfrak{J} is bijective. Hence we get the following proposition.

Proposition 2. *The semigroup $\mathbf{End}_{\langle \sigma \rangle}(\mathcal{C}_+(a, b))$ is isomorphic to the direct product $\mathfrak{RZ}(\omega) \times (\mathbb{N}, *)$.*

Fix an arbitrary $b^i a^j \in \mathcal{C}_+(a, b)$, $i, j \in \omega, j \geq i$. Then for any $\sigma_{l, m} \in \mathbf{End}_{\langle \sigma \rangle}(\mathcal{C}_+(a, b))$ and any $\lambda_k \in \mathbf{End}_{\langle \lambda \rangle}(\mathcal{C}_+(a, b))$ we have that

$$\begin{aligned} ((b^i a^j) \sigma_{l, m}) \lambda_k &= \begin{cases} (1) \lambda_k, & \text{if } i = j; \\ ((b^l a^{l+m})^{j-i}) \lambda_k, & \text{if } i < j \end{cases} = \\ &= \begin{cases} 1, & \text{if } i = j; \\ (b^l a^{l+m(j-i)}) \lambda_k, & \text{if } i < j \end{cases} = \\ &= \begin{cases} 1, & \text{if } i = j; \\ b^{kl} a^{kl+km(j-i)}, & \text{if } i < j \end{cases} = \\ &= (b^i a^j) \sigma_{kl, km} \end{aligned}$$

and

$$\begin{aligned} ((b^i a^j)\lambda_k)\sigma_{l,m} &= (b^{ki} a^{kj})\sigma_{l,m} = \\ &= \begin{cases} (b^{ki} a^{ki})\sigma_{l,m}, & \text{if } ki = kj; \\ (b^l a^{l+m})^{kj-ki}, & \text{if } ki < kj \end{cases} = \\ &= \begin{cases} (b^{ki} a^{ki})\sigma_{l,m}, & \text{if } i = j; \\ (b^l a^{l+m})^{kj-ki}, & \text{if } i < j \end{cases} = \\ &= \begin{cases} 1, & \text{if } i = j; \\ b^l a^{l+km(j-i)}, & \text{if } i < j \end{cases} = \\ &= (b^i a^j)\sigma_{l,km}. \end{aligned}$$

This implies the following

Proposition 3. $\sigma_{l,m}\lambda_k = \sigma_{kl,km}$ and $\lambda_k\sigma_{l,m} = \sigma_{l,km}$ for any $\sigma_{l,m} \in \mathbf{End}_{\langle\sigma\rangle}(\mathcal{C}_+(a, b))$ and any $\lambda_k \in \mathbf{End}_{\langle\lambda\rangle}(\mathcal{C}_+(a, b)) \setminus \{\lambda_0\}$.

By $\mathbf{End}_{\langle\lambda,\sigma\rangle}(\mathcal{C}_+(a, b))$ we denote the subsemigroup of $\mathbf{End}(\mathcal{C}_+(a, b))$, which is generated by the families $\{\lambda_k : k \in \omega\}$ and $\{\sigma_{l,m} : l, m \in \omega, m > 0\}$ of endomorphisms of the monoid $\mathcal{C}_+(a, b)$. We summarise the results of this section in the following theorem.

Theorem 1. (1) λ_1 is the identity element of the semigroup $\mathbf{End}(\mathcal{C}_+(a, b))$, and hence it is the identity element of the semigroups $\mathbf{End}_{\langle\lambda\rangle}(\mathcal{C}_+(a, b))$ and $\mathbf{End}_{\langle\lambda,\sigma\rangle}(\mathcal{C}_+(a, b))$;

(2) λ_0 is the zero of the semigroup $\mathbf{End}(\mathcal{C}_+(a, b))$, and hence it is the zero of $\mathbf{End}_{\langle\lambda\rangle}(\mathcal{C}_+(a, b))$ and $\mathbf{End}_{\langle\lambda,\sigma\rangle}(\mathcal{C}_+(a, b))$;

(3) the set $I = \mathbf{End}_{\langle\sigma\rangle}(\mathcal{C}_+(a, b)) \cup \{\lambda_0\}$ is an ideal of the semigroup $\mathbf{End}_{\langle\lambda,\sigma\rangle}(\mathcal{C}_+(a, b))$.

Proof. Statements (1) and (2) are trivial.

(3) By Proposition 3 we have that $\sigma_{l,m}\lambda_k, \lambda_k\sigma_{l,m} \in \mathbf{End}_{\langle\sigma\rangle}(\mathcal{C}_+(a, b))$ for any $\sigma_{l,m} \in \mathbf{End}_{\langle\sigma\rangle}(\mathcal{C}_+(a, b))$ and $\lambda_k \in \mathbf{End}_{\langle\lambda\rangle}(\mathcal{C}_+(a, b)) \setminus \{\lambda_0\}$. Since λ_0 is the zero of the semigroup $\mathbf{End}(\mathcal{C}_+(a, b))$, the above arguments imply that I is an ideal of the semigroup $\mathbf{End}_{\langle\lambda,\sigma\rangle}(\mathcal{C}_+(a, b))$. \square

2. On monoid injective endomorphisms of $\mathcal{C}_+(a, b)$

Example 1. For arbitrary positive integer n and arbitrary $s=0, \dots, n-1$ we define the mapping $\lambda_{n,s} : \mathcal{C}_+(a, b) \rightarrow \mathcal{C}_+(a, b)$ by the formula

$$(b^i a^j)\lambda_{n,s} = \begin{cases} a^{nj}, & \text{if } i = 0; \\ b^{ni-s} a^{nj-s}, & \text{if } i \neq 0 \end{cases}$$

for all $i, j \in \omega$.

Proposition 4. *For any positive integer n and any $s = 0, \dots, n - 1$ the map $\lambda_{n,s}$ is an injective monoid endomorphism of the monoid $\mathcal{C}_+(a, b)$.*

Proof. Fix any positive integers i, j, k, l such that $i \leq j$ and $k \leq l$, and non-negative integers m and q . Then we have that

$$\begin{aligned} (b^i a^j \cdot b^k a^l) \lambda_{n,s} &= \begin{cases} (b^{i-j+k} a^l) \lambda_{n,s}, & \text{if } j < k; \\ (b^i a^l) \lambda_{n,s}, & \text{if } j = k; \\ (b^i a^{j-k+l}) \lambda_{n,s}, & \text{if } j > k \end{cases} = \\ &= \begin{cases} b^{n(i-j+k)-s} a^{nl-s}, & \text{if } j < k; \\ b^{ni-s} a^{nl-s}, & \text{if } j = k; \\ b^{ni-s} a^{n(j-k+l)-s}, & \text{if } j > k, \end{cases} \end{aligned}$$

$$\begin{aligned} (b^i a^j) \lambda_{n,s} \cdot (b^k a^l) \lambda_{n,s} &= b^{ni-s} a^{nj-s} \cdot b^{nk-s} a^{nl-s} = \\ &= \begin{cases} b^{(ni-s)-(nj-s)+(nk-s)} a^{nl-s}, & \text{if } nj - s < nk - s; \\ b^{ni-s} a^{nl-s}, & \text{if } nj - s = nk - s; \\ b^{ni-s} a^{nj-s-(nk-s)+(nl-s)}, & \text{if } nj - s > nk - s \end{cases} = \\ &= \begin{cases} b^{n(i-j+k)-s} a^{nl-s}, & \text{if } j < k; \\ b^{ni-s} a^{nl-s}, & \text{if } j = k; \\ b^{ni-s} a^{n(j-k+l)-s}, & \text{if } j > k, \end{cases} \end{aligned}$$

$$\begin{aligned} (b^i a^j \cdot a^m) \lambda_{n,s} &= (b^i a^{j+m}) \lambda_{n,s} = \\ &= b^{ni-s} a^{n(j+m)-s} = \\ &= b^{ni-s} a^{nj-s} \cdot a^{mn} = \\ &= (b^i a^j) \lambda_{n,s} \cdot (a^m) \lambda_{n,s}, \end{aligned}$$

$$\begin{aligned} (a^m \cdot b^i a^j) \lambda_{n,s} &= \begin{cases} (b^{i-m} a^j) \lambda_{n,s}, & \text{if } m < i; \\ (a^{m-i+j}) \lambda_{n,s}, & \text{if } m \geq i \end{cases} = \\ &= \begin{cases} b^{n(i-m)-s} a^{nj-s}, & \text{if } m < i; \\ a^{n(m-i+j)}, & \text{if } m \geq i, \end{cases} \end{aligned}$$

$$\begin{aligned} (a^m) \lambda_{n,s} \cdot (b^i a^j) \lambda_{n,s} &= a^{mn} \cdot b^{ni-s} a^{nj-s} = \\ &= \begin{cases} b^{ni-s-nm} a^{nj-s}, & \text{if } mn < ni - s; \\ a^{mn-(ni-s)+nj-s}, & \text{if } mn \geq ni - s \end{cases} = \end{aligned}$$

$$\begin{aligned}
 &= \begin{cases} b^{n(i-m)-s} a^{nj-s}, & \text{if } m < i - s/n; \\ a^{n(m-i+j)}, & \text{if } m \geq i - s/n \end{cases} = \\
 &= \begin{cases} b^{n(i-m)-s} a^{nj-s}, & \text{if } m < i; \\ a^{n(m-i+j)}, & \text{if } m \geq i, \end{cases}
 \end{aligned}$$

because $s = 0, \dots, n - 1$, and

$$(a^m \cdot a^q)\lambda_{n,s} = (a^{m+q})\lambda_{n,s} = a^{(m+q)n} = a^{mn} \cdot a^{qn} = (a^m)\lambda_{n,s} \cdot (a^q)\lambda_{n,s}.$$

Hence the map $\lambda_{n,s}$ is a monoid endomorphism. The condition that $s = 0, \dots, n - 1$ implies that $\lambda_{n,s}$ is an injective map. \square

By $\mathbf{End}_{\langle \lambda^\infty \rangle}(\mathcal{C}_+(a, b))$ we denote the subset of $\mathbf{End}(\mathcal{C}_+(a, b))$, which consists of the elements of the family $\{\lambda_{n,s} : n \in \omega, s = 0, \dots, n - 1\}$ of endomorphisms of the monoid $\mathcal{C}_+(a, b)$.

Let S and T be arbitrary semigroups. Let $\varphi : T \rightarrow \mathbf{End}(S)$, $t \mapsto \varphi_t$ be a homomorphism from T into the semigroup $\mathbf{End}(S)$ of endomorphisms of S . The *semidirect product* of S and T defined on the product $S \times T$ with the semigroup operation

$$(s_1, t_1) \cdot (s_2, t_2) = (s_1 \cdot (s_2)\varphi_{t_1}, t_1 \cdot t_2),$$

and it is denoted by $S \rtimes_\varphi T$.

Theorem 2. *The set $\mathbf{End}_{\langle \lambda^\infty \rangle}(\mathcal{C}_+(a, b))$ is a submonoid of $\mathbf{End}(\mathcal{C}_+(a, b))$ and $\mathbf{End}_{\langle \lambda^\infty \rangle}(\mathcal{C}_+(a, b))$ is isomorphic to a submonoid of the semidirect product $(\mathbb{N}, *) \rtimes_\varphi (\omega, +)$, where $(p)\varphi_n = np$.*

Proof. Fix arbitrary positive integers n_1, n_2 , $s_1 = 0, \dots, n_1 - 1$ and $s_2 = 0, \dots, n_2 - 1$. Then for any $b^i a^j \in \mathcal{C}_+(a, b)$ we have that

$$\begin{aligned}
 ((b^i a^j)\lambda_{n_1, s_1})\lambda_{n_2, s_2} &= \begin{cases} (a^{n_1 j})\lambda_{n_2, s_2}, & \text{if } i = 0; \\ (b^{n_1 i - s_1} a^{n_1 j - s_1})\lambda_{n_2, s_2}, & \text{if } i \neq 0 \end{cases} = \\
 &= \begin{cases} a^{n_1 n_2 j}, & \text{if } i = 0; \\ a^{n_2(n_1 j - s_1)}, & \text{if } i \neq 0 \text{ and } n_1 i - s_1 = 0; \\ b^{n_2(n_1 i - s_1) - s_2} a^{n_2(n_1 j - s_1) - s_2}, & \text{if } i \neq 0 \text{ and } n_1 i - s_1 \neq 0 \end{cases} = \\
 &= \begin{cases} a^{n_1 n_2 j}, & \text{if } i = 0; \\ b^{n_2 n_1 i - (s_1 n_2 + s_2)} a^{n_2 n_1 j - (s_1 n_2 + s_2)}, & \text{if } i \neq 0 \end{cases} = \\
 &= (b^i a^j)\lambda_{n_1 n_2, s_1 n_2 + s_2}.
 \end{aligned}$$

Since $s_1 < n_1$ and $s_2 < n_2$, we have that

$$s_1 n_2 + s_2 \leq (n_1 - 1)n_2 + s_2 < (n_1 - 1)n_2 + n_2 = n_1 n_2 - n_2 + n_2 = n_1 n_2,$$

and hence $s_1n_2 + s_2 < n_1n_2$. This implies that $\mathbf{End}_{\langle \lambda^\infty \rangle}(\mathcal{C}_+(a, b))$ is a subsemigroup of $\mathbf{End}(\mathcal{C}_+(a, b))$. Since $\lambda_{n,0} = \lambda_n$ for any positive integer n , $\mathbf{End}_{\langle \lambda \rangle}(\mathcal{C}_+(a, b))$ is a submonoid of $\mathbf{End}_{\langle \lambda^\infty \rangle}(\mathcal{C}_+(a, b))$.

We define the map $\Phi: \mathbf{End}_{\langle \lambda^\infty \rangle}(\mathcal{C}_+(a, b)) \rightarrow (\mathbb{N}, *) \rtimes_{\varphi} (\omega, +)$ by the formula $(\lambda_{n,s})\Phi = (n, s)$. The above arguments imply that

$$\begin{aligned} (\lambda_{n_1,s_1} \lambda_{n_2,s_2})\Phi &= (\lambda_{n_1n_2,s_1n_2+s_2})\Phi = \\ &= (n_1n_2, s_1n_2 + s_2) = \\ &= (n_1n_2, (s_1)\varphi_{n_2} + s_2) = \\ &= (n_1, s_1)(n_2, s_2) = \\ &= (\lambda_{n_1,s_1})\Phi(\lambda_{n_2,s_2})\Phi, \end{aligned}$$

and hence Φ is a homomorphism. □

Example 2. We define the map $\varsigma: \mathcal{C}_+(a, b) \rightarrow \mathcal{C}_+(a, b)$ by the formula

$$(b^i a^j)\varsigma = \begin{cases} 1, & \text{if } i = j = 0; \\ b^{i+1} a^{j+1}, & \text{otherwise} \end{cases}$$

for any $i, j \in \omega$.

Lemma 3. *The map $\varsigma: \mathcal{C}_+(a, b) \rightarrow \mathcal{C}_+(a, b)$ is an injective monoid endomorphism.*

Proof. A simple verification shows that ς is an injective map. Obviously that it is sufficient to show that for any $b^i a^j \neq 1$ and $b^k a^l \neq 1$ the following equality $(b^i a^j \cdot b^k a^l)\varsigma = (b^i a^j)\varsigma \cdot (b^k a^l)\varsigma$ holds. Indeed,

$$\begin{aligned} (b^i a^j \cdot b^k a^l)\varsigma &= \begin{cases} (b^{i-j+k} a^l)\varsigma, & \text{if } j < k; \\ (b^i a^l)\varsigma, & \text{if } j = k; \\ (b^i a^{j-k+l})\varsigma, & \text{if } j > k \end{cases} = \\ &= \begin{cases} b^{i-j+k+1} a^{l+1}, & \text{if } j < k; \\ b^{i+1} a^{l+1}, & \text{if } j = k; \\ b^{i+1} a^{j-k+l+1}, & \text{if } j > k \end{cases} \end{aligned}$$

and

$$(b^i a^j)\varsigma \cdot (b^k a^l)\varsigma = b^{i+1} a^{j+1} \cdot b^{k+1} a^{l+1} =$$

$$\begin{aligned}
 &= \begin{cases} b^{i+1-(j+1)+k+1}a^{l+1}, & \text{if } j + 1 < k + 1; \\ b^{i+1}a^{l+1}, & \text{if } j + 1 = k + 1; \\ b^{i+1}a^{j+1-(k+1)+l+1}, & \text{if } j + 1 > k + 1 \end{cases} = \\
 &= \begin{cases} b^{i-j+k+1}a^{l+1}, & \text{if } j < k; \\ b^{i+1}a^{l+1}, & \text{if } j = k; \\ b^{i+1}a^{j-k+l+1}, & \text{if } j > k, \end{cases}
 \end{aligned}$$

and hence ς is an injective monoid endomorphism of $\mathcal{C}_+(a, b)$. □

By $\mathbf{End}_{\langle \varsigma \rangle}(\mathcal{C}_+(a, b))$ we denote the subsemigroup of $\mathbf{End}(\mathcal{C}_+(a, b))$, which is generated by endomorphism ς of the monoid $\mathcal{C}_+(a, b)$. Also by $\mathbf{End}_{\langle \varsigma \rangle}^1(\mathcal{C}_+(a, b))$ we denote the semigroup $\mathbf{End}_{\langle \varsigma \rangle}(\mathcal{C}_+(a, b))$ with the adjoined unit. Without loss of generality we may assume that

$$\mathbf{End}_{\langle \varsigma \rangle}^1(\mathcal{C}_+(a, b)) = \mathbf{End}_{\langle \varsigma \rangle}(\mathcal{C}_+(a, b)) \cup \{\lambda_1\}.$$

Proposition 5. *The semigroup $\mathbf{End}_{\langle \varsigma \rangle}(\mathcal{C}_+(a, b))$ is isomorphic to the additive semigroup of positive integers $(\mathbb{N}, +)$, and hence $\mathbf{End}_{\langle \varsigma \rangle}^1(\mathcal{C}_+(a, b))$ is isomorphic to the additive monoid of non-negative integers $(\omega, +)$.*

Proof. For any $b^i a^j \in \mathcal{C}_+(a, b)$ and any positive integer n by the definition of ς we have that

$$\begin{aligned}
 (b^i a^j)\varsigma^n &= ((b^i a^j)\varsigma)\varsigma^{n-1} = \\
 &= \begin{cases} (1)\varsigma^{n-1}, & \text{if } i = j = 0; \\ (b^{i+1}a^{j+1})\varsigma^{n-1}, & \text{otherwise} \end{cases} = \\
 &= \dots = \\
 &= \begin{cases} (1)\varsigma, & \text{if } i = j = 0; \\ (b^{i+n-1}a^{j+n-1})\varsigma, & \text{otherwise} \end{cases} = \\
 &= \begin{cases} 1, & \text{if } i = j = 0; \\ b^{i+n}a^{j+n}, & \text{otherwise.} \end{cases}
 \end{aligned}$$

The definition of the bicyclic monoid $\mathcal{C}(a, b)$ implies that $b^{k_1}a^{l_1} = b^{k_2}a^{l_2}$ in $\mathcal{C}_+(a, b)$ if and only if $k_1 = k_2$ and $l_1 = l_2$. This and above equalities imply that the endomorphism ς generates the infinite cyclic subsemigroup in $\mathbf{End}(\mathcal{C}_+(a, b))$, and hence $\mathbf{End}_{\langle \varsigma \rangle}(\mathcal{C}_+(a, b))$ is isomorphic to the additive semigroup of positive integers $(\mathbb{N}, +)$. The last statement of the proposition is obvious. □

Lemma 4. *Let $\varepsilon: \mathcal{C}_+(a, b) \rightarrow \mathcal{C}_+(a, b)$ be an injective monoid endomorphism such that $(a)\varepsilon = a^n$ for some positive integer n . Then there exists $s \in \{0, \dots, n - 1\}$ such that $\varepsilon = \lambda_{n,s}$.*

Proof. We observe that if ε is an injective endomorphism of $\mathcal{C}_+(a, b)$ then for any idempotents $b^i a^i, b^j a^j \in \mathcal{C}_+(a, b)$ the inequality $b^i a^i \preceq b^j a^j$ implies that $(b^i a^i)\varepsilon \preceq (b^j a^j)\varepsilon$, because the equality $b^i a^i \cdot b^j a^j = b^i a^i$ implies that

$$(b^i a^i)\varepsilon \cdot (b^j a^j)\varepsilon = (b^i a^i \cdot b^j a^j)\varepsilon = (b^i a^i)\varepsilon.$$

Since ε is an injective monoid endomorphism of $\mathcal{C}_+(a, b)$, we conclude that $(b^i a^i)\varepsilon \neq (b^j a^j)\varepsilon$ and $(b^0 a^0)\varepsilon = (1)\varepsilon = 1 = b^0 a^0$. Hence there exists a strictly increasing sequence $\{s_i\}_{i \in \omega}$ in ω such that $(b^i a^i)\varepsilon = b^{s_i} a^{s_i}$ for any $i \in \omega$ and $s_0 = 0$. Then for any positive integer i we have that

$$\begin{aligned} (b^i a^{i+1})\varepsilon &= (b^i a^i \cdot a)\varepsilon = \\ &= (b^i a^i)\varepsilon \cdot (a)\varepsilon = \\ &= b^{s_i} a^{s_i} \cdot a^n = \\ &= b^{s_i} a^{s_i+n} \end{aligned}$$

and

$$\begin{aligned} (b^i a^{i+1})\varepsilon &= (a \cdot b^{i+1} a^{i+1})\varepsilon = \\ &= (a)\varepsilon \cdot (b^{i+1} a^{i+1})\varepsilon = \\ &= a^n \cdot b^{s_{i+1}} a^{s_{i+1}} = \\ &= \begin{cases} a^n, & \text{if } n \geq s_{i+1}; \\ b^{s_{i+1}-n} a^{s_{i+1}}, & \text{if } n < s_{i+1}. \end{cases} \end{aligned}$$

The injectivity of ε implies that $n < s_{i+1}$. Since $s_0 = 0$ and $s_i < s_{i+1}$ for any $i \in \omega$, the above equalities imply that $s_{i+1} = s_i + n$ for any positive integer i , and hence $s_{i+1} - s_i = n$ for $i \in \mathbb{N}$. By induction we have that $s_{i+1} = s_1 + in$ for all $i \in \mathbb{N}$. This implies that

$$(b^i a^i)\varepsilon = b^{(i-1)n+s_1} a^{(i-1)n+s_1}$$

for any positive integer i . Then we get that

$$\begin{aligned} (b^i a^{i+l})\varepsilon &= (b^i a^i \cdot a^l)\varepsilon = \\ &= (b^i a^i)\varepsilon \cdot (a^l)\varepsilon = \\ &= b^{(i-1)n+s_1} a^{(i-1)n+s_1} \cdot ((a)\varepsilon)^l = \\ &= b^{(i-1)n+s_1} a^{(i-1)n+s_1} \cdot (a^n)^l = \\ &= b^{(i-1)n+s_1} a^{(i-1)n+s_1} \cdot a^{nl} = \\ &= b^{(i-1)n+s_1} a^{(i+l-1)n+s_1} \end{aligned}$$

for any $i, l \in \omega$. Also, since

$$\begin{aligned} a^n &= (a)\varepsilon = \\ &= (a \cdot ba)\varepsilon = \\ &= (a)\varepsilon \cdot (ba)\varepsilon = \\ &= a^n \cdot b^{s_1} a^{s_1}, \end{aligned}$$

properties of semigroup operation of $\mathcal{C}_+(a, b)$ and the natural partial order on the set of idempotents of the monoid $\mathcal{C}_+(a, b)$ imply that $s_1 \leq n$, and by injectivity of ε we have that $s_1 > 0$. Put $s = n - s_1$. Then we obtain that $s \in \{0, \dots, n - 1\}$ and

$$\begin{aligned} (b^i a^{i+l})\varepsilon &= b^{(i-1)n+s_1} a^{(i+l-1)n+s_1} = \\ &= b^{in-n+s_1} a^{(i+l)n-n+s_1} = \\ &= b^{in-s} a^{(i+l)n-s}. \end{aligned}$$

This and Proposition 4 imply that $\varepsilon = \lambda_{n,s}$. □

Proposition 6. *Let S, T , and U be semigroups, let $\mathfrak{h}: S \rightarrow T$ be a homomorphism, and let $\mathfrak{g}: U \rightarrow T$ be an injective homomorphism. If $(S)\mathfrak{h} \subseteq (U)\mathfrak{g}$, then the mapping $\mathfrak{f}: S \rightarrow U$ which is defined by the formula $(s)\mathfrak{f} = ((s)\mathfrak{h})\mathfrak{g}^{-1}$ is a homomorphism. Moreover, if the homomorphism $\mathfrak{h}: S \rightarrow T$ is injective or a monoid homomorphism, then so is \mathfrak{f} , too.*

Proof. Since $\mathfrak{g}: U \rightarrow T$ is an injective homomorphism, the map $\mathfrak{f}: S \rightarrow U$ is well defined. Also, for arbitrary $s_1, s_2 \in S$ we have that

$$\begin{aligned} (s_1 \cdot s_2)\mathfrak{f} &= ((s_1 \cdot s_2)\mathfrak{h})\mathfrak{g}^{-1} = \\ &= ((s_1)\mathfrak{h} \cdot (s_2)\mathfrak{h})\mathfrak{g}^{-1} = \\ &= ((s_1)\mathfrak{h})\mathfrak{g}^{-1} \cdot ((s_2)\mathfrak{h})\mathfrak{g}^{-1} = \\ &= (s_1)\mathfrak{f} \cdot (s_2)\mathfrak{f}, \end{aligned}$$

because $\mathfrak{g}: U \rightarrow T$ is an injective homomorphism. Hence $\mathfrak{f}: S \rightarrow U$ is a homomorphism. The second statement of the proposition is obvious. □

Lemma 5. *Let $\varepsilon: \mathcal{C}_+(a, b) \rightarrow \mathcal{C}_+(a, b)$ be an injective monoid endomorphism such that $(a)\varepsilon = b^n a^{n+p}$ for some positive integers n and p . Then there exists $s \in \{0, \dots, n - 1\}$ such that $\varepsilon = \lambda_{p,s} \varepsilon^n$.*

Proof. Since ε is an injective monoid endomorphism of $\mathcal{C}_+(a, b)$, arguments presented in the proof of Lemma 4 imply that there exists a strictly increasing sequence $\{s_i\}_{i \in \omega}$ in ω such that $(b^i a^i)\varepsilon = b^{s_i} a^{s_i}$ for any $i \in \omega$ and $s_0 = 0$. Then for any positive integers i and j we have that

$$\begin{aligned} (b^i a^{i+j})\varepsilon &= (b^i a^i \cdot a^j)\varepsilon = \\ &= (b^i a^i)\varepsilon \cdot (a^j)\varepsilon = \\ &= b^{s_i} a^{s_i} \cdot (b^n a^{n+p})^j = \\ &= b^{s_i} a^{s_i} \cdot b^n a^{n+pj} = \\ &= \begin{cases} b^n a^{n+pj}, & \text{if } n \geq s_i; \\ b^{s_i} a^{s_i+pj}, & \text{if } n < s_i. \end{cases} \end{aligned}$$

This implies that $(\mathcal{C}_+(a, b))\varepsilon \subseteq (\mathcal{C}_+(a, b))\zeta^n$. By Proposition 6 the mapping $f: \mathcal{C}_+(a, b) \rightarrow \mathcal{C}_+(a, b)$ which is defined by the formula $(b^i a^j)f = ((b^i a^j)\varepsilon)(\zeta^n)^{-1}$ is an endomorphism of the monoid $\mathcal{C}_+(a, b)$. Simple verifications show that $(a)f = a^p$. By Lemma 4 we have that $f = \lambda_{p,s}$ for some $s \in \{0, \dots, n - 1\}$, i.e., $\lambda_{p,s} = \varepsilon(\zeta^n)^{-1}$. Since ζ is an injective monoid endomorphism of $\mathcal{C}_+(a, b)$, we conclude that $\varepsilon = \lambda_{p,s}\zeta^n$. \square

Theorem 3 describes all injective endomorphisms of the semigroup $\mathcal{C}_+(a, b)$ and it follows from Lemmas 4 and 5.

Theorem 3. *Let $\varepsilon: \mathcal{C}_+(a, b) \rightarrow \mathcal{C}_+(a, b)$ be an injective monoid endomorphism. Then only one of the following statements holds:*

- (1) *there exist a positive integer p and $s \in \{0, \dots, p - 1\}$ such that $\varepsilon = \lambda_{p,s}$;*
- (2) *there exist positive integers n, p , and $s \in \{0, \dots, p - 1\}$ such that $\varepsilon = \lambda_{p,s}\zeta^n$.*

It is natural to ask the following: what is a semigroup operation on the subsemigroup $\mathbf{End}_{\langle \lambda\zeta \rangle}(\mathcal{C}_+(a, b))$ of $\mathbf{End}(\mathcal{C}_+(a, b))$ of endomorphisms of $\mathcal{C}_+(a, b)$ which is generated by endomorphism of the form $\lambda_{p,s}\zeta^n$, where $p, n \in \mathbb{N}$ and $s \in \{0, \dots, n - 1\}$. Theorem 4 describes the structure of the semigroup operation on the semigroup $\mathbf{End}_{\langle \lambda\zeta \rangle}(\mathcal{C}_+(a, b))$ of injective monoid endomorphisms of $\mathcal{C}_+(a, b)$.

Theorem 4. *$\mathbf{End}_{\langle \lambda\zeta \rangle}(\mathcal{C}_+(a, b))$ is a subsemigroup of $\mathbf{End}(\mathcal{C}_+(a, b))$ and it is isomorphic to the subsemigroup of the Cartesian power \mathbb{N}^3 with the following semigroup operation*

$$(p_1, s_1, n_1) \cdot (p_2, s_2, n_2) = (p_2 p_1, p_2 s_1, p_2 n_1 - s_2 + n_2).$$

Proof. Fix arbitrary $p_1, n_1, p_2, n_2 \in \mathbb{N}$, $s_1 \in \{0, \dots, n_1 - 1\}$ and $s_2 \in \{0, \dots, n_2 - 1\}$. Then for any $b^i a^j \in \mathcal{C}_+(a, b)$ we have that

$$\begin{aligned} & (((b^i a^j) \lambda_{p_1, s_1}) \zeta^{n_1}) \lambda_{p_2, s_2} \zeta^{n_2} = \\ & = \begin{cases} ((a^{p_1 j}) \zeta^{n_1}) \lambda_{p_2, s_2} \zeta^{n_2}, & \text{if } i = 0; \\ (((b^{p_1 i - s_1} a^{p_1 j - s_1}) \zeta^{n_1}) \lambda_{p_2, s_2}) \zeta^{n_2}, & \text{if } i \neq 0 \end{cases} = \\ & = \begin{cases} ((1) \lambda_{p_2, s_2}) \zeta^{n_2}, & \text{if } i = 0 \text{ and } j = 0; \\ ((b^{n_1} a^{p_1 j + n_1}) \lambda_{p_2, s_2}) \zeta^{n_2}, & \text{if } i = 0 \text{ and } j \neq 0; \\ (((b^{p_1 i - s_1 + n_1} a^{p_1 j - s_1 + n_1}) \lambda_{p_2, s_2}) \zeta^{n_2}), & \text{if } i \neq 0 \end{cases} = \\ & = \begin{cases} (1) \zeta^{n_2}, & \text{if } i = 0 \text{ and } j = 0; \\ (b^{p_2 n_1 - s_2} a^{p_2(p_1 j + n_1) - s_2}) \zeta^{n_2}, & \text{if } i = 0 \text{ and } j \neq 0; \\ (b^{p_2(p_1 i - s_1 + n_1) - s_2} a^{p_2(p_1 j - s_1 + n_1) - s_2}) \zeta^{n_2}, & \text{if } i \neq 0 \end{cases} = \\ & = \begin{cases} 1, & \text{if } i = 0 \text{ and } j = 0; \\ b^{p_2 n_1 - s_2 + n_2} a^{p_2(p_1 j + n_1) - s_2 + n_2}, & \text{if } i = 0 \text{ and } j \neq 0; \\ b^{p_2(p_1 i - s_1 + n_1) - s_2 + n_2} a^{p_2(p_1 j - s_1 + n_1) - s_2 + n_2}, & \text{if } i \neq 0 \end{cases} = \\ & = \begin{cases} 1, & \text{if } i = 0 \text{ and } j = 0; \\ b^{p_2 n_1 - s_2 + n_2} a^{p_2 p_1 j + p_2 n_1 - s_2 + n_2}, & \text{if } i = 0 \text{ and } j \neq 0; \\ b^{p_2 p_1 i - p_2 s_1 + p_2 n_1 - s_2 + n_2} a^{p_2 p_1 j - p_2 s_1 + p_2 n_1 - s_2 + n_2}, & \text{if } i \neq 0 \end{cases} \\ & = (b^i a^j) \lambda_{p_2 p_1, p_2 s_1} \zeta^{p_2 n_1 - s_2 + n_2} \end{aligned}$$

The above equalities imply the statement of the theorem. □

Remark 1. 1. The endomorphisms λ_k , $k \in \omega$, $\lambda_{p, s}$, $p, n \in \mathbb{N}$, $s \in \{0, \dots, n - 1\}$ and ζ of the monoid $\mathcal{C}_-(a, b)$ are introduced by the same formulae.

2. Repeating the proofs of corresponding statements in Sections 1 and 2 we obtain that the same statements about the corresponding endomorphisms of the monoid $\mathcal{C}_-(a, b)$. Moreover, the corresponding semigroups of endomorphisms are pairwise isomorphic.

Acknowledgements

The authors acknowledge Alex Ravsky and the referee for they comments and suggestions.

References

- [1] Clifford, A.H., Preston, G.B.: The Algebraic Theory of Semigroups. Vol. I., Amer. Math. Soc. Surveys 7, Providence, R.I. (1961)
- [2] Clifford, A.H., Preston, G.B.: The Algebraic Theory of Semigroups. Vol. II., Amer. Math. Soc. Surveys 7, Providence, R.I. (1967)
- [3] Descalço, L., Ruškuc, N.: Subsemigroups of the bicyclic monoid. *Int. J. Algebra Comput.* **15**(1), 37–57 (2005). <https://doi.org/10.1142/S0218196705002098>
- [4] Descalço, L., Ruškuc, N.: Properties of the subsemigroups of the bicyclic monoid. *Czech. Math. J.* **58**(2), 311–330 (2008). <https://doi.org/10.1007/s10587-008-0018-7>
- [5] Gutik, O.: On non-topologizable semigroups. Preprint at <https://arxiv.org/abs/2405.16992> (2024)
- [6] Gutik, O., Prokhorenkova, O., Sekh, D.: On endomorphisms of the bicyclic semigroup and the extended bicyclic semigroup. *Visn. L'viv. Univ., Ser. Mekh.-Mat.* **92**, 5–16 (2021) (in Ukrainian). <https://doi.org/10.30970/vmm.2021.92.005-016>
- [7] Lawson, M.: Inverse Semigroups. The Theory of Partial Symmetries. World Scientific, Singapore (1998). <https://doi.org/10.1142/3645>
- [8] Makanjuola, S.O., Umar, A.: On a certain subsemigroup of the bicyclic semigroup. *Commun. Algebra.* **25**(2), 509–519 (1997). <https://doi.org/10.1080/00927879708825870>
- [9] Wagner, V.V.: Generalized groups. *Dokl. Akad. Nauk SSSR.* **84**, 1119–1122 (1952) (in Russian)

CONTACT INFORMATION

O. Gutik,
Sh.-A. Penza

Ivan Franko National University of Lviv,
Universytetska 1, Lviv, 79000, Ukraine
E-Mail: oleg.gutik@lnu.edu.ua,
sher-ali.penza@lnu.edu.ua

Received by the editors: 22.08.2024
and in final form 16.11.2024.