

The structure of automorphism groups of semigroup inflations

Ganna Kudryavtseva

Communicated by B. V. Novikov

ABSTRACT. In this paper we prove that the automorphism group of a semigroup being an inflation of its proper subsemigroup decomposes into a semidirect product of two groups one of which is a direct sum of full symmetric groups.

1. Introduction

In the study of a specific semigroup the description of all its automorphisms is one of the most important questions. The automorphism groups of many important specific semigroups are described (see, for example, [7] and references therein). It can be observed that for two types of semigroups of rather diverse nature the automorphism groups have similarities in their structure: each of them decomposes into a semidirect product of two groups one of which being a direct sum of full symmetric groups. The mentioned two types of semigroups are variants of some semigroups of mappings (see [4], [6]) and maximal nilpotent subsemigroups of some transformation semigroups (see [2], [5]).

In the present paper we establish a general result, which, in particular, unifies all the above-mentioned examples. We deal with an abstract semigroup being an inflation of its proper subsemigroup, constructed as follows. Let S be a semigroup. First introduce an equivalence relation h on S via $(a, b) \in h$ if and only if $ax = bx$ and $xa = xb$ for all $x \in S$, that is a and b are not distinguished by multiplication from either side.

2000 Mathematics Subject Classification: 20M10.

Key words and phrases: *semigroup inflation, semidirect product, automorphism group.*

We would like to remark that the history of the study of h stems back at least to [6], where it was considered for the variants of certain semigroups of mappings. Set further

$$\psi = (h \cap ((S \setminus S^2) \times (S \setminus S^2))) \cup \{(a, a) : a \in S^2\}.$$

Denote by T any transversal of ψ . Then T is a subsemigroup of S , and S is an inflation of T . An automorphism τ of T will be called *extendable* provided that τ coincides with the restriction to T of a certain automorphism of S . Clearly, all extendable automorphisms of T constitute a subgroup, H , of the group $\text{Aut}T$ of all automorphisms of T .

We state our result as the following theorem.

Theorem 1. *The group $\text{Aut}S$ is isomorphic to a semidirect product of two groups one of which (the one which is normal) is the direct sum of the full symmetric groups on the ψ -classes and the other one is the group H consisting of all extendable automorphisms of T .*

All the papers [2], [4], [5], [6] deal precisely with the "semigroup inflation construction" introduced in this paper (though this construction is not mentioned explicitly in any of these papers), and their main results concerning automorphism groups can be deduced from our present general result — Theorem 1. Recently Theorem 1 was applied to describe automorphism groups of certain partition semigroups (see [3], Section 11 for details).

2. Construction

A semigroup S is called an *inflation* of its subsemigroup (see [1], Section 3.2) T provided that there is an onto map $\theta : S \rightarrow T$ such that:

- $\theta^2 = \theta$;
- $a\theta b\theta = ab$ for all $a, b \in S$.

In the described situation S is often referred to as an *inflation of T with an associated map θ* (or just *with a map θ*).

It is immediate that if S is an inflation of T then T is a retract of S (that is the image under an idempotent homomorphism) and that $S^2 \subset T$.

Lemma 2. *Suppose that S is an inflation of T with the map θ . Then $\ker\theta \subset h$.*

Proof. Let $(a, b) \in \ker\theta$ and $s \in S$. Then

$$as = a\theta s\theta = b\theta s\theta = bs; sa = s\theta a\theta = s\theta b\theta = sb,$$

which implies that $(a, b) \in h$. \square

Lemma 3. *The equivalence ψ , defined in the Introduction, is a congruence on S .*

Proof. Obviously, ψ is an equivalence relation. Prove that ψ is left and right compatible. Let $(a, b) \in \psi$ and $a \neq b$. Then $(a, b) \in (h \cap ((S \setminus S^2) \times (S \setminus S^2)))$. Let $c \in S$. Since $(a, b) \in h$ we have that $ac = bc$ and $ca = cb$ for each $c \in S$. It follows that $(ac, bc) \in \psi$ and $(ca, cb) \in \psi$ as ψ is reflexive. \square

Fix an arbitrary transversal of ψ and denote it by T .

Lemma 4. *T is a subsemigroup of S , and S is an inflation of T .*

Proof. T is a subsemigroup of S as $T \supset S^2$. Let θ be the map $S \rightarrow T$ which sends any element x from S to the unique element of the ψ -class of x , belonging to T . The construction implies that S is an inflation of T with the map θ . \square

Let $S = \cup_{a \in T} X_a$ be a decomposition of S into the union of ψ -classes, where X_a denotes the ψ -class of a . Set G_a to be the full symmetric group acting on X_a and $G = \oplus_{a \in T} G_a$.

Lemma 5. *π is an automorphism of S , for every $\pi \in G$.*

Proof. It is enough to show that $(xy)\pi = x\pi y\pi$ whenever $x, y \in S$. Suppose first that $x, y \in S \setminus S^2$. Since $xy \in S^2$ it follows that π stabilizes xy , so that $(xy)\pi = xy$. Now, the inclusions $(x, x\pi) \in h$ and $(y, y\pi) \in h$ imply $x\pi y\pi = x\pi y = xy$. This yields $xy\pi = x\pi y\pi$, and the proof is complete. \square

The following proposition gives a characterization of extendable automorphisms of T .

Proposition 6. *An automorphism τ of T is extendable if and only if the following condition holds:*

$$(\forall a, b \in T) \quad a\tau = b \Rightarrow |X_a| = |X_b|. \quad (1)$$

Proof. Suppose $\tau \in \text{Aut}T$ is extendable and $a \in T$. In the case when $a \in S \setminus S^2$ we have

$$X_a = \{b \in S \mid (a, b) \in h \text{ and } b \in S \setminus S^2\}.$$

Clearly, $(a, b) \in h \iff (a\tau, b\tau) \in h$ and $b \in S \setminus S^2 \iff b\tau \in S \setminus S^2$ for all $a, b \in S$. It follows that

$$X_{a\tau} = \{b\tau \mid b \in X_a\},$$

which implies (1). The inclusion $a \in S^2$ is equivalent to $a\tau \in S^2$. But then $|X_a| = |X_{a\tau}| = 1$, which also implies (1).

Suppose now that (1) holds for certain $\tau \in \text{Aut}T$. Then one can extend τ to $\bar{\tau} \in \text{Aut}S$ as follows.

Fix a collection of sets $I_a, a \in T$, and bijections $f_a : I_a \rightarrow X_a, a \in T$, satisfying the following conditions:

- $|I_a| = |X_a|$;
- $I_a = I_b$ whenever $|X_a| = |X_b|$;
- $I_a \cap I_b = \emptyset$ whenever $|X_a| \neq |X_b|$;
- if $a, b \in T$ and $|X_a| = |X_b|$ then $af_a^{-1} = bf_b^{-1}$.

It is straightforward that such collections $I_a, a \in T$, and $f_a, a \in T$, exist.

Consider $x \in S \setminus T$. Since T is a transversal of ψ , there is $a \in T$ such that $x \in X_a$. By the hypothesis we have $|X_a| = |X_{a\tau}|$. Set $\bar{\tau}$ on X_a to be the map from X_a to $X_{a\tau}$ defined via $x \mapsto xf_a^{-1}f_{a\tau}$. In this way we define a permutation $\bar{\tau}$ of S such that $\bar{\tau}|_T = \tau$. It will be called an *extension* of τ to S . To complete the proof, we are left to show that $\bar{\tau}$ is a homomorphism. Let $x, y \in S, x \in X_a, y \in X_b$. Then

$$(xy)\bar{\tau} = (ab)\tau = a\tau b\tau = x\bar{\tau}y\bar{\tau},$$

as required. □

Let $\tau \in H$. Of course, $\bar{\tau}$, constructed in the proof of Proposition 6, depends not only on τ , but also on the sets I_a and the maps f_a , so that τ may have several extensions to S . Fix any extension $\bar{\tau}$ of τ .

Lemma 7. $\tau \mapsto \bar{\tau}$ is an embedding of H into $\text{Aut}S$.

Proof. Proof is immediate by the construction of $\bar{\tau}$. □

Denote by \bar{H} the image of H under the embedding of H into $\text{Aut}S$ from Lemma 7.

3. Proof of Theorem 1

Proposition 8. \overline{H} acts on G by automorphisms via $\pi^\tau = \tau^{-1}\pi\tau$, $\tau \in \overline{H}$, $\pi \in G$.

Proof. Let $\pi \in G$ and $\tau \in \overline{H}$. Show first that $\pi^\tau \in G$. Take any $x \in S$. Let X_a be the block which contains x . We consequently have $x\tau^{-1} \in X_{a\tau^{-1}}$, $x\tau^{-1}\pi \in X_{a\tau^{-1}}$ and $x\tau^{-1}\pi\tau \in X_{a\tau^{-1}\tau} = X_a$. Hence $x\pi^\tau \in X_a$. It follows that $\pi^\tau \in G$.

That $\pi \mapsto \pi^\tau$ is one-to-one, onto and homomorphic immediately follows from its definition. We are left to show that the map sending $\tau \in \overline{H}$ to $\pi \mapsto \pi^\tau \in \text{Aut}G$ is homomorphic. The latter follows from the equalities $\pi^{\tau_1\tau_2} = (\tau_1\tau_2)^{-1}\pi(\tau_1\tau_2) = (\pi^{\tau_1})^{\tau_2}$. The proof is complete. \square

In the following two lemmas we show that G and \overline{H} intersect by the identity automorphism and generate $\text{Aut}S$.

Lemma 9. $G \cap \overline{H} = \{id\}$, where id is the identity automorphism of S .

Proof. The proof follows from the observation that the decomposition $S = \cup_{a \in T} X_a$ is fixed by each element of G , while only by the identity element of \overline{H} . \square

Lemma 10. $\text{Aut}S = \overline{H} \cdot G$.

Proof. Let $\varphi \in \text{Aut}S$. The definition of ψ implies that φ maps each ψ -class onto some other ψ -class. Define a bijection $\tau : T \rightarrow T$ via $a\tau = b$ provided that $X_a\varphi = X_b$. Show that τ is an extendable automorphism of T . The definition of τ assures that (1) holds, and thereby τ is extendable in view of Proposition 6. Let $\overline{\tau} \in \text{Aut}S$ be an extension of τ . The construction implies that $\varphi(\overline{\tau})^{-1} \in G$. \square

Now the proof of Theorem 1 follows from Proposition 8 and Lemmas 9 and 10.

4. Acknowledgements

The author expresses her gratitude to Prof. B.V. Novikov for the careful reading of the initial version of this paper.

References

- [1] A.H. Clifford, G.B. Preston, *The algebraic theory of semigroups*, Amer. Math. Soc. Surveys 7 (Providence, R.I.) 1961, 1967.
- [2] O. Ganyushkin, S. Temnikov and H. Shafranov, *Automorphism groups of maximal nilpotent subsemigroups of the semigroup $\mathcal{IS}(M)$* , Math. Stud., 13 (2000), 1, 11-22.

- [3] G. Kudryavtseva and V. Maltcev, *On the structure of two generalizations of the full inverse symmetric semigroup*, math.GR/0602623.
- [4] G. Kudryavtseva and G. Tsyaputa, *The automorphism group of the sandwich inverse symmetric semigroup*, Visnyk of Kyiv University, 13-14 (2005), 101-105 (in Ukrainian), English translation: math.GR/0509678.
- [5] A. Stronska, *On the automorphisms for the nilpotent subsemigroups of the order-decreasing transformation semigroup*, Proceedings of the F. Skorina Gomel University, 2 (2007), 38-49.
- [6] J.S.V. Symons, *On a generalization of the transformation semigroup*, J. Austral Math. Soc., 19 (1975), 47-61.
- [7] F. Szechtman, *On the automorphism group of the centralizer of an idempotent in the full transformation monoid*, Semigroup Forum, 70 (2005), 238-242.

CONTACT INFORMATION

G. Kudryavtseva Department of Mechanics and Mathematics,
Kyiv Taras Shevchenko University,
Volodymyrs'ka, 60, Kyiv, 01033, Ukraine
E-Mail: akudr@univ.kiev.ua

Received by the editors: 19.11.2005
and in final form 28.05.2007.